

Short Note

2-Methyl-7-(phenylsulfanylmethyl)naphthalene

Jane Bogdanov ^{†,*} and Przemyslaw Maslak

Department of Chemistry, The Pennsylvania State University, University Park, Pennsylvania 16802, USA; E-Mail: nyl@psu.edu

[†] Present address: Institute of Chemistry, Faculty of Natural Sciences & Mathematics, Sts. Cyril and Methodius University, PO Box 162, 1000 Skopje, Macedonia.

* Author to whom correspondence should be addressed; E-Mail: j_b_bogdanov@yahoo.com; Tel.: +389 2 324 9918; Fax: +389 2 3226865.


Received: 16 March 2010 / Accepted: 24 March 2010 / Published: 30 March 2010

Abstract: A new sulfide, 2-methyl-7-(phenylsulfanylmethyl)naphthalene was synthesized and its MS, IR, ¹H NMR, ¹³C NMR and DEPT-135 data are reported.

Keywords: phase-transfer catalysis; 2-methyl-7-(phenylsulfanylmethyl)naphthalene; naphthylmethyl phenyl sulfide

1. Introduction

In connection with our investigation of competitive mesolytic cleavages [1] of radical anions, we needed 2-methyl-7-(phenylsulfanylmethyl)naphthalene (**3**) as gas chromatography standard. The synthetic method of choice for this naphthylmethyl phenyl sulfide is reaction between naphthylmethyl halide and a corresponding thiophenoxide. From the various developed procedures [2–7] we have decided to utilize the phase-transfer variant used by Guthrie and Maslak [7] for preparation of the analogous ethers. The starting material for these derivatives was 2-bromomethyl-7-methylnaphthalene (**1**), which could be conveniently obtained by benzylic bromination of the commercially available 2,7-dimethylnaphthalene, using the procedure described by Buuhoi and co-workers [8]. We would like to report a convenient procedure for preparation of 2-methyl-7-(phenylsulfanylmethyl)naphthalene.

Scheme 1. Synthesis of 2-methyl-7-(phenylsulfanylmethyl)naphthalene.

2. Experimental

2.1. General

All chemicals were obtained from commercial sources and were used without further purification. Melting points were determined using a Mel-Temp apparatus and are uncorrected. NMR spectra were recorded on a Bruker 400 MHz instrument, using deuteriochloroform as solvent and tetramethylsilane as internal standard. The IR spectra were recorded on a Perkin Elmer Model 1600 instrument between sodium chloride plates in carbon tetrachloride. The number of hydrogens on each carbon was determined from ¹³C NMR and DEPT-135 spectra. The mass spectra were recorded on a Kratos MS-25 RFA double-focusing mass spectrometer in electron impact (EI) mode. Gas chromatography was performed on a Varian 3700 instrument with packed column. The column was 1/8" in diameter and 50 cm in length packed with 5% OV-101 on supelcoport, purchased from Supelco. The carrier gas was helium (30 mL/min flow), the detection was accomplished with flame ionization and monitored with HP-3390A reporting integrator. TLC was carried out using Merck pre-coated plates (60 F₂₅₄, 250 μm).

2.2. 2-Methyl-7-(phenylsulfanylmethyl)naphthalene (3)

A mixture of 2-bromomethyl-7-methylnaphthalene (1) (0.200 g, 0.861 mmol), 2:1 (v/v) dichloromethane/acetonitrile (5 mL), 1M NaOH (5 mL), freshly distilled thiophenol (0.188 g, 1.71 mmol) and a drop of methyltricaprylammonium chloride (MTCAC) was vigorously stirred at 60 °C for 3 h under argon. The mixture was cooled to room temperature, the layers were separated and the organic layer was washed with 1M sodium hydroxide (2 × 10 mL), water (2 × 10 mL) and brine (1 × 10 mL). Drying over sodium sulfate and removal of the solvent *in vacuo* afforded a yellow solid. Recrystallization from ethanol yielded 0.121 g (54%) of 3 as white plates.

M.p. 89–91 °C.

R_f (20% dichloromethane in hexanes) = 0.26.

GC: R_t = 15.80 min (100 °C, 3 min, 8 °C/min to 280 °C).

^1H NMR (400 MHz, CDCl_3): δ = 7.72 (d, J = 8.4 Hz, 1 H), 7.69 (d, J = 8.4 Hz, 1 H), 7.58 (s, 1 H), 7.50 (s, 1 H), 7.38 (dd, J = 8.4 Hz, 1.7 Hz, 1 H), 7.34–7.29 (m, 2 H), 7.27 (dd, J = 8.4 Hz, 1.7 Hz, 1 H), 7.24–7.12 (m, 3 H), 4.24 (s, 2 H, Ar- CH_2 -SPh), 2.48 (s, 3 H, Ar- CH_3).

^{13}C NMR (100 MHz): δ = 136.35 (C), 135.78 (C), 134.87 (C), 133.52 (C), 130.82 (C), 129.90 (CH), 128.83 (CH), 128.08 (CH), 127.99 (CH), 127.43 (CH), 126.78 (CH), 126.68 (CH), 126.35 (CH), 126.08 (CH) (Ar), 39.41 (Ar- CH_2 -SPh), 21.70 (Ar- CH_3).

DEPT-135 NMR: δ = 129.90 (\uparrow , CH), 128.83 (\uparrow , CH), 128.08 (\uparrow , CH), 127.99 (\uparrow , CH), 127.43 (\uparrow , CH), 126.78 (\uparrow , CH), 126.68 (\uparrow , CH), 126.35 (\uparrow , CH), 126.08 (\uparrow , CH) (Ar), 39.41 (\downarrow , Ar- CH_2 -SPh), 21.70 (\uparrow , Ar- CH_3).

IR (CCl_4 , cm^{-1}): 3055, 3022, 2924, 2861, 1638, 1610, 1585, 1515, 1480, 1439, 1383, 1336, 1230, 1078, 1025, 1012, 959, 902.

EI-MS (m/z , rel. intensity): 265 ($\text{M}^+ + 1$, 3%), 264 (M^+ , 7%), 197 (3%), 157 (9%), 156 (9%), 155 ($\text{M}^+ - \text{SPh}$, 100%), 111 (14%).

3. Conclusion

The reaction of 2-bromomethyl-7-methylnaphthalene with thiophenol under phase-transfer catalytic conditions, in the presence of methyltricaprylammonium chloride (MTCAC) gave the desired product **3** in moderate yield. The structure of **3** was unambiguously determined by spectroscopic means (^1H NMR, ^{13}C NMR, DEPT-135 NMR, IR and MS). The reaction can be conveniently monitored by GC and/or TLC (20% dichloromethane in hexanes). The reaction was performed in degassed solution under argon atmosphere to avoid the oxidation of the thiophenol into diphenyl disulfide [9]. Diphenyl disulfide can pose a separation problem and can interfere with the mesolytic cleavage studies.

References

1. Maslak, P.; Narvaez, J.N. Mesolytic cleavage of C-C bonds. Comparison with homolytic and heterolytic processes in the same substrate. *Angew. Chem. Int. Ed. Engl.* **1990**, *29*, 283–285.
2. Frankel, M.; Gertner, D.; Jacobson, H.; Zilkha, A. Syntheses of Poly-S-Alkyl-L-Cysteines. *J. Chem. Soc.* **1960**, 1390–1393.
3. Dymicky, M.; Byler, D.M. Synthesis of ethyl *n*-carbobenzoxytyrosyl-*s*-benzylcysteinylglycinate. *Org. Prep. Proced. Int.* **1991**, *23*, 93–101.
4. Vögtle, F.; Klieser, B. Cesium-ion assisted synthesis of strained compounds. *Synthesis* **1982**, 294–296.
5. Yin, J.M.; Pidgeon, C. A simple and efficient method for preparation of unsymmetrical sulfides. *Tetrahedron Lett.* **1997**, *38*, 5953–5954.
6. Herriott, A.W.; Picker, D. Phase-transfer synthesis of sulfides and dithioacetals. *Synthesis* **1975**, 447–448.
7. Maslak, P.; Guthrie, R.D. Electron apportionment in cleavage of radical-anions. 2. naphthylmethyl phenyl ethers vs. naphthyl benzyl ethers. *J. Am. Chem. Soc.* **1986**, *108*, 2637–2640.

8. Buuhoi, N.P.; Lecocq, J. Side-chain bromination of some alkylnaphthalenes with *n*-bromosuccinimide. *J. Chem. Soc.* **1946**, 830–832.
9. McMurry, J. *Organic Chemistry*, 5th ed.; Brooks/Cole: Pacific Grove, CA, USA, 2000; p. 729.

© 2010 by the authors; licensee Molecular Diversity Preservation International, Basel, Switzerland. This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).

Copyright of Molbank is the property of Molecular Diversity Preservation International (MDPI) and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.