

## AMIDE AND THIOAMIDE BANDS OF BENZANILIDE AND THIOBENZANILIDE IN THE VIBRATIONAL SPECTRA

I. PETROV and O. GRUPEĆ

Faculty of Chemistry, Cyril and Methodius University,  
Skopje, Yugoslavia

### ABSTRACT

It has been widely accepted that in secondary amides and secondary thioamides, in the spectral region between 1600 and 1200  $\text{cm}^{-1}$  two characteristic bands could be recognized (Amide II and Amide III for amides and B and C bands for thioamides). Our spectra of benzanilide,  $\text{C}_6\text{H}_5(\text{C}=\text{O})\text{NHC}_6\text{H}_5$ , and thiobenzanilide,  $\text{C}_6\text{H}_5(\text{C}=\text{S})\text{NHC}_6\text{H}_5$ , show that in this region there are, at least, four prominent bands which shift on deuteration. That could indicate that all these bands are in connection with the vibrations of amide and/or thioamide groups. Some other amide and thioamide bands have been also discussed.

### INTRODUCTION

The infrared spectra of amides and thioamides have been a subject of investigation of many authors (ref. 1-9). The assignments of the characteristic bands have been made, both empirically and by the use of normal coordinate calculations with various degree of sophistication.

The infrared spectrum of thiobenzanilide has been already reported and some assignments have been made (ref. 3,4). Our study (ref. 10) of the N-H stretching region in the infrared spectra of thiobenzanilide showed that thiobenzanilide crystallizes in four different modifications, which we designated as  $\alpha$ ,  $\beta$ ,  $\gamma$  and  $\delta$  phases. The structural differences among these phases are related mainly to the differences in the hydrogen bonding (ref. 10).

This paper is mainly concerned with infrared and Raman spectra of benzanilide and thiobenzanilide\*, in the region below 1600  $\text{cm}^{-1}$ . We examined this spectral region in hope that it might extend our

---

\*In this article, we will discuss only  $\alpha$  phase, since similar behavior has been noticed in the spectra of the other thiobenzanilide polymorphs.

understanding of the nature of the characteristic amide and thioamide bands.

#### EXPERIMENTAL

Benzanilide was prepared from aniline and benzoylchloride and than by action of  $P_2S_5$  converted to thiobenzanilide.

Benzanilide and thiobenzanilide were deuterated by dissolving them in dioxan, adding  $D_2O$  until precipitation occurred and drying the separated crystals in vacuo.

The infrared spectra were recorded mainly as KBr discs on a Perkin-Elmer 580 IR Spectrophotometer and Raman spectra on a Jeol JRS-S1B Laser Raman Spectrophotometer. For recording the infrared spectra at liquid nitrogen temperature (LNT), the variable VLT-2 cell (RIIC) was used.

#### RESULTS AND DISCUSSIONS

In the infrared spectra of benzanilide and thiobenzanilide, it is expected that two bands should disappear in the region between  $1600$  and  $1200\text{ cm}^{-1}$ , on deuteration. Instead of that, four prominent bands disappear. At the same time, in deuterated spectra of examined compounds, instead of one, two strong bands appear, in the region between  $1500$  and  $1400\text{ cm}^{-1}$  ( see Fig. 1 and 2).




Fig. 1. Infrared spectrum of (a) protonated and (b) deuterated benzanilide

In the spectrum of D-benzanilide (see Fig. 1), in the lower frequency range, another band of medium intensity at  $938\text{ cm}^{-1}$  can be seen, while in D-thiobenzanilide (see Fig. 2) there are three new bands in the same region. In the Raman spectra (see Fig. 3 and 4), the bands that occur in the region between  $1600$  and  $1400\text{ cm}^{-1}$  are weak while the bands in the region between  $1400$  and  $1200\text{ cm}^{-1}$  are quite strong, in both, benzanilide and thiobenzanilide.

A possible explanation for appearance of four appreciably strong bands in the range between 1600 and 1400  $\text{cm}^{-1}$ , which shift on deuteration and have different intensities in the Raman and infrared




Fig. 2. Infrared spectrum of (a) protonated and (b) deuterated thiobenzanilide




Fig. 3. Raman spectrum of benz-anilide

which in most cases couples with N-H vibration. If this is so, then it should be expected that this band is rather strong in the Raman spectrum of thiobenzanilide. However, in our Raman spectrum this band appears as a very weak shoulder on the low frequency side of the very strong breathing band of the benzene ring, at 1000  $\text{cm}^{-1}$  (see Fig. 4). We believe that the appearance of this band is mainly due to a contribution of  $\nu\text{C}=\text{S}$ . In deuterated thiobenzanilide  $\delta\text{N}-\text{D}$  mode probably couples extensively with this vibration to give rise to three new bands at 1250, 1070 and 940  $\text{cm}^{-1}$  (see Fig. 2).

spectra, may be that these bands are in connection with symmetrical and asymmetrical vibrational modes in which two non-equivalent C-N groups are involved. These modes probably couple strongly with  $\delta\text{N}-\text{H}$  mode giving rise to four bands. Thus, two strong bands which appear

between 1500 and 1400  $\text{cm}^{-1}$  in the spectra of deuterated compounds, may be due to the two stretching vibrations of the two nonequivalent C-N groups.

A strong and sharp band at 985  $\text{cm}^{-1}$ , in the infrared spectra of thiobenzanilide is considered to be localized  $\nu\text{C}=\text{S}$  vibration (ref. 3,4,9). However, Jensen and Nielsen (ref. 7) designated it as D band with major contribution of symmetrical stretching NCS vibration,




Fig. 4. Raman spectrum of thio-benzanilide

band almost certainly corresponds to  $\gamma$ C-H of the phenyl ring. The  $690\text{ cm}^{-1}$  band can also be assigned to phenyl bending mode. Therefore only the band at  $710\text{ cm}^{-1}$ , which appear in the infrared spectra of both, benzanilide and thiobenzanilide, may partly belong to a vibration of the amide and/or thioamide group.

#### ACKNOWLEDGEMENT

The financial support of the Community for Scientific Activities of the Socialist Republic of Macedonia is gratefully acknowledged.

#### REFERENCES

- 1 R. Fraser and W. Prace, *Nature* **170**, (1959), 490 pp; *Proc. Roy Soc.* **141B**, (1953), 66 pp.
- 2 T. Miyazawa, T. Shimanouchi and S. Mizushima, *J. Chem. Phys.* **29**, (1958), 611 pp.
- 3 D. Hadži, *J. Chem. Soc.* (1957), 849 pp.
- 4 C. N. Rao and R. Venkataraghavan, *Spectrochim. Acta* **18**, (1962), 541 pp.
- 5 C. Collard-Charon and M. Renson, *Bull. Soc. Chim. Belges* **72**, (1963), 304 pp.
- 6 H. O. Desseyne and M. A. Herman, *Spectrochim. Acta* **23A**, (1967), 2457 pp.
- 7 K. A. Jensen and P. H. Nielsen, *Acta Chim. Scand.* **20**, (1966), 597pp
- 8 C. N. R. Rao and G. C. Chaturvedi, *Spectrochim. Acta* **27A**, (1971), 520 pp.
- 9 H. E. Hallam and C. M. Jones, *Spectrochim. Acta* **25A**, (1969), 1791 pp.
- 10 O. Grupče and I. Petrov, to be published.

The infrared spectrum of benzanilide in the region below  $900\text{ cm}^{-1}$  is very similar to that of the thio-benzanilide. In the range between  $760$  and  $600\text{ cm}^{-1}$ , three strong bands could be seen in both of the examined compounds (see Fig. 1 and 2). Jensen and Nielsen (ref. 7) considered that  $767/752$  and  $710\text{ cm}^{-1}$  bands of thiobenzanilide are due to mixed vibrations of the phenyl, C=S and N-H groups. We believe that the  $760\text{ cm}^{-1}$