

WEEDS AND WEED MANAGEMENT IN CONSERVATION AGRICULTURE (CA) INFLUENCED BY ITS THREE PILLARS AND HERBICIDES[#]

Zvonko Pacanoski

Faculty for Agricultural Sciences and Food, University Ss. Cyril and Methodius,
Skopje, Republic of Macedonia

e-mail: zvonkop@fznh.ukim.edu

CA is based on three large pillars: reducing or eliminating tilling, covering the soil, diversifying crop rotations. One of the most difficult management issues within this system are weeds and proper weed management. CA brings a shift in dynamics of the weed seed bank, weed populations, density and growth, requiring clear understanding and formulating a strategy for their management. Crop residues influence weed seed germination and seedling emergence by interfering with sunlight availability and suppression through physical and chemical allelopathic effects. Cover crops are fundamental and sustainable tools to manage weeds reducing their numbers physically and chemically. Diversification of the crop rotation help to disrupt the growing cycle of weeds, and prevent any weed species to dominate. Herbicides are an integral part of weed management in CA, as well. When herbicides are not used at appropriate rates or in rotation caused environmental pollution, weed shift, and resistance development in some weeds.

Key words: CA; weeds; crop residues; cover crops; crop rotation

REASONS FOR CA APPEARING AND ITS PILLARS

Conventional tillage (CT) has been an integral part of crop production since crops were first cultivated. Some historians have even evaluated the progress of agrarian societies by their developments in tillage. The premise for CT was to subdue or destroy native vegetation so desired plants might develop free from competition. From the onset of agriculture, CT was synonymous with seedbed preparation and weed control [1]. Apart this, CT is used for crop establishment in order to loosen and aerate the soil for planting, incorporate crop residues and nutrients, enhance the release of nutrients from the soil for crop growth, regulate the circulation of water and air within the soil [2, 3] and when carried out in autumn/winter expose the soil to frost in order to benefit soil structure [4].

Nevertheless CT practices often increase soil erosion rates leading to deteriorating soil physical, chemical and biological properties [5], reduced soil

quality such as soil structure, with consequences for water infiltration [6], poor soil porosity, nutrient loss and low organic matter content [7], lead to increased greenhouse gas emissions [8] and create hard pans below the plough layer, as well [9]. Poor soil nutrient statuses in combination with poor weed management practices often contribute to decreased yields [10]. To alleviate this challenge, researchers have suggested a more sustainable method of farming, commonly referred to as Conservation Agriculture (CA).

CA (synonymous of zero tillage farming or no-tillage farming, ridge-till, mulch-till, and noninversion tillage) emerged historically as a response to soil erosion crises in the USA, Brazil, Argentina and Australia where currently, it spans over million hectares. The vulnerability of plough-based agriculture was exposed during the Dust Bowl era (1931-39); as the wind blew away the precious top soil from the drought-ravaged southern plains of the US, leaving behind the failed crops and farms. However, there was no answer then to solve the question of soil

[#]Dedicated to academician Gjorgji Filipovski on the occasion of his 100th birthday

degradation. Then, what *Nature* magazine termed "an agricultural bombshell" was dropped by Edward Faulkner on July 5, 1943; with the first issue of his masterpiece book "*Plowman's Folly and A Second Look*" [11]. This book was a milestone in the history of agricultural practices—he questioned the wisdom of ploughing. Some of his statements are: "No one has ever advanced a scientific reason for plowing"; "There is simply no need for plowing in the first instance. And most of the operations that customarily follow the plowing are entirely unnecessary, if the land has not been plowed"; "There is nothing wrong with our soil, except our interference"; and, "It can be said with considerable truth that the use of the plow has actually destroyed the productiveness of our soils." The statements were questioned by both farmers and researchers because alternatives to ploughing at that time did not allow farmers to control weeds or plant into the residues [12]. The idea was widely embraced by farmers all across the world, particularly in the USA only after the Second World War, when the development of chemicals for agriculture allowed them to try it out. With the introduction of 2,4-D in the mid-1940s, producers were, for the first time, given an economical chemical alternative to tillage for preplant weed control. The introduction of numerous other herbicides in the succeeding decades allowed reduced and CA systems to become more feasible and popular [13]. CA systems are being advocated since the 1970s. However, the majority of CA expansion worldwide has occurred since the mid- to late-1990s [14]. This has been accelerated due to the development of efficient farm machinery and the availability of effective herbicides coupled with trained manpower, which have resulted in reduced production costs and higher profitability, besides several indirect benefits [15].

CA is a system designed to achieve agricultural sustainability by improving the biological functions of the agroecosystem with limited mechanical practices and judicious use of chemical inputs. According to FAO [16], CA is an approach to managing agro-ecosystems for improved and sustained productivity, increased profits and food security while preserving and enhancing the resource base and the environment. CA is characterized by three linked principles, namely:

1. Continuous no- or minimal mechanical soil disturbance (i.e., no-tillage and direct sowing or broadcasting of crop seeds, and direct placing of planting material in the soil; minimum soil disturbance from cultivation, harvest operation or farm traffic, in special cases limited strip tillage);

2. Permanent organic soil cover, especially by crop residues, crops and cover crops to protect and feed the soil, develop surface mulch; and

3. Diversification of crop species grown in sequence or associations through rotations or, in case of perennial crops, associations of plants, including a balanced mix of legume and non legume crops to help moderate possible weed, disease and pest problems, generate biomass, fix atmospheric nitrogen and serve as nutrient pumps [17–19].

These three principles converge towards one central and stated goal: to reduce soil degradation and improve soil fertility, by preserving its organic matter, flora and fauna [20].

CA, particularly no-till (NT) system as the core principle of CA, offers a way of optimizing productivity and ecosystem services, offering a wide range of economic, environmental and social benefits to the producer and to the society. At the same time, no-till farming is enabling agriculture to respond to some of the global challenges associated with climate change, land and environmental degradation, and increasing cost of food, energy and production inputs [21–23].

BENEFITS AND LIMITATIONS OF CA

CA technologies are essentially herbicide-driven, machine-driven and knowledge-driven, and therefore require vastly-improved expertise and resources for adoption in large areas. Like every technology, CA mainly is characterized as beneficiary oriented, but it is faced with many challenges and limitations, as well.

A host of benefits can be achieved through employing components of CA, including reduced soil erosion (96 % less erosion) [24] and water runoff (reduction of 70 % in the volume of run-off) [25, 26], increased productivity through improved soil quality, soil water holding capacity increased [27], water availability, water infiltration and water use efficiency [21], reduced sediment and fertilizer pollution in lakes and streams, increased biotic diversity, reduced labour demands [18, 28, 29] and improved quality of life. Advantages also include climate change mitigation through reduced emissions due to 60–70 % lower fuel use, 20–50 % lower fertilizer and pesticides use, 50 % reduction in machinery and labour requirement [30], C-sequestration $0.85 \text{ t ha}^{-1} \text{ y}^{-1}$ or more [31], and no CO_2 release as a result of no burning of residues. Overall, CA has a much lower carbon footprint than CT agriculture [32], and greenhouse gas (GHG) emissions of CO_2 , CH_4 and NO_2 are all reduced with CA [33–35]. Based on experiences from North and South Ameri-

ca, increased soil C sequestration is reported as the chief reason for improved soil productivity with time under CA compared to CT fields [36]. Improved soil moisture conservation [6], reduced erosion, and increased biological activity [37], are all associated with residue retention. Crop rotation increases crop diversity, reducing crop yield penalties associated with insect damage, diseases and weed infestations whilst improving nutrient cycling [38].

Despite both environmental and production advantages offered through CA, adoption rates have previously lagged in many countries due to several factors including: availability of required equipment, lack of information, producer mindsets, and, initially, greater weed control problems, herbicide dependency [38, 39] and increased risk of herbicide resistance [40]. Other disadvantages include reduced spring soil temperature [41], increased pest problems (incidence of rhizoctonia disease, for example), risk of increased N₂O emissions and increased dissolved reactive P leaching, reduced reliability of crop yields, risk of topsoil compaction [42], and reluctance of farmers to accept the new practice [43]. Several factors including biophysical, socio-economic and cultural limit the adoption of CA, particularly by resource-poor farmers. The current major barriers to the spread of CA systems are (i) competing use of crop residues in rainfed areas, (ii) weed management strategies, particularly for perennial species, (iii) localized insect and disease infestation, and (iv) likelihood of lower crop productivity if site-specific component technologies are not adopted [44, 45]

WEEDS IN CA SYSTEMS

Although benefits are context specific, CA has been identified as an effective tool for sustainably increasing yields in many parts of the world [21, 46], adopting CA will face several managerial changes, and weed control is perceived as one of the most challenging [5, 47–49].

CT has been a major agricultural weed control technique for several decades [50], so the development of CA systems that advocate NT or reduced tillage has significant implications for growers [51]. CT affects weeds by uprooting, dismembering, and burying them deep enough to prevent emergence, by changing the soil environment and so promoting or inhibiting the weeds' germination and establishment, and by moving their seeds both vertically and horizontally [52–54]. CT is also used to incorporate herbicides into the soil and to remove surface residues that might otherwise impede the herbicides' effectiveness. Reduction in tillage intensity and fre-

quency, as practiced under CA, generally increases weed infestation and causes a variation in the dynamics of the weed populations [55].

Mishra and Singh [56] observed that over the course of time, an NT–NT sequence favored relatively higher weed growth over a CT–CT sequence in a rice–wheat system. While weed growth in the initial year was not higher under the NT–NT sequence, in the third year of experimentation total weed dry weight was significantly higher under the NT–NT than CT–CT tillage sequence. Total weed density was significantly lower (16.3 plants m⁻²) under the CT than the other reduced tillage systems (36.7–39.2 plants m⁻²). The main benefit of CT is a highly significant decline of perennial weeds. Only 2.6 perennial weed plants per quadrant in CT as compared to 7.5–9.0 in reduced tillage treatments were noted [57]. Tolimir *et al.* [58] also noted considerably lower weed infestation per square meter under CT (7 weeds) compared to reduced (39 weeds) and NT (46 weeds). Swanton *et al.* [59] found that tillage was an important factor affecting weed composition: *Chenopodium album* and *Amaranthus retroflexus* were associated with a CT system, whereas *Digitaria sanguinalis* was associated with NT.

Shifts in weed populations from annual large-seeded broadleaf to annuals grass and small-seeded broadleaf and perennials, as well have been observed in CA systems [60–63]. Perennial weeds thrive in reduced or NT systems [64, 65]. Most perennial weeds have the ability to reproduce from several structural organs other than seeds. Among them, perennial monocots are considered a greater threat than perennial dicots in the adoption of reduced tillage systems [11]. In practice, researchers have shown that small-seeded annual grass species and perennials become more difficult to manage as tillage is decreased, whereas large-seeded broadleaf weeds become easier to manage in production systems with less tillage [66]. For example, in ZT-DSR (zero-tillage- dry direct-seeding), weed flora often shifts towards more difficult to control and competitive grasses and sedges such as *Leptochloa chinensis* (L.) Nees, *Eragrostis* spp., *Echinochloa colona* (L.) Link., *Cyperus* spp., *Eclipta prostrata* (L.) L., *Ammannia* spp., *Sphenochloa zeylinica* Gaertn. [67]. Similar, in the Eastern Indo-Gangetic Plains, big and serious problems under NT are *Cyperus rotundus* L. and *Cynodon dactylon* (L.) Pers. [68]. According to Shaw *et al.* [1] winter annual and biennial weeds, and some brush species that were not problematic with CT can increase with NT, like *Cirsium arvense*, where NT leaves the roots undisturbed and *Cirsium arvense* populations can increase. Opposite, analysis of multiplied dominance ratio (MDR) of

weeds in Japan showed that summer annual weeds, especially grass weeds, were much more abundant than perennial weeds in the NT fields [69].

Different tillage systems disturb the vertical distribution of weed seeds in the soil, in different ways. Studies have found that moldboard plowing buries most weed seeds in the tillage layer, whereas chisel plowing leaves most of the weed seeds closer to the soil surface [70]. In reduced or NT systems, depending on the soil type, 60–90 % of weed seeds are located in the top 5 cm of the soil [52, 54, 71]. Similar, the highest number of weed seed species was found in the treatments with reduced and NT treatments in a soil layer of 0–5 cm. In deeper soil layers (5–10, 10–20 cm), no differences in weed seed species number were found [72]. Chauhan *et al.* [73] reported that a low-soil-disturbance single-disc system retained more than 75 % of the weed seeds in the top 1-cm soil layer, whereas the high-soil-disturbance seeding system buried more than 75 % of the seeds to a depth more than 5 cm. Further, the accumulation of the weed seeds at the soil surface increases their chance to germinate in one season with suitable moisture and temperature, and they are exposed to insect predation, (vertebrates and invertebrates) [74] fungal and bacterial attack [75], rodents and birds consume [76] and decay thus depletion of the weed seed bank is high [77] as well as weed density [78]. Sagar and Mortimer [79] found that weed seed survival across time is lowest when seeds remain on the soil surface because of exposure to environmental extremes and predation. Similar, Usman *et al.* [80] concluded that there is a rapid loss of viability of weed seeds in addition to predation on the soil surface in NT compared to CT where seeds are buried in the soil and prevented from environmental hazards. Even in the absence of seed predators, weed seeds from species as diverse as *Panicum miliaceum* L., *Amaranthus retroflexus* L., and *Solanum sarrachoides* Sendtner have been shown to lose viability at a greater rate when positioned near the soil surface than when buried below the emergence zone [81, 82].

Although some studies found that weed seed bank, weed density and weed infestation are smaller in NT compared with CT, other studies claimed the opposite [83]. According to Singh *et al.* [11] the presence of weed seeds on the upper soil surface, due to no tillage operation, leads to higher weed infestation in the NT system. Cardina *et al.* [63] studied the weed seed bank size and composition after 35 years of continuous crop rotation and tillage system and concluded that weed seed density was highest in NT and generally declined as tillage intensity increased. The soil weed seed bank was 1.5 and 2.2

times greater in the shallow ploughing and shallow ploughless tillage treatments, compared with the CT treatment (deep ploughing). In the shallow ploughing and shallow ploughless tillage treatments, there were found 25.5 % and 41.5 % more weed seed species in the soil, compared with the CT treatment [84]. According to Menalled *et al.* [85], above ground weed biomass, species density, and diversity were lowest in the CT system, intermediate in the NT system, and highest in the low-input and organic systems. Higher weed seed densities in NT systems may be the result of reduced herbicides availability because of adsorption to near surface organic matter [86].

COVER CROPS AND THEIR RESIDUES

The use of cover crops in CA offers many advantages, one of which is weed suppression through physical as well as chemical allelopathic effects [87, 88], while actively growing or after termination [89]. Prior to termination, cover crops can compete with weed species for necessary resources such as light, water, and nutrients; cover crops can also release allelochemicals into the soil which may be detrimental to nearby competing weed species, particularly for small-seeded weeds [90]. Cereal rye (*Secale cereale* L.) and soft red winter wheat (*Triticum aestivum* L.) used as cover crops also contain allelopathic compounds that inhibit weed growth [91]. Yenish *et al.* [92] reported increased short-term weed control using a rye cover crop in NT corn (*Zea mays* L.) but not season-long control. In southern Brazil, black oat (*Avena strigosa* Schreb.) is the predominant cover crop on millions of hectares of NT soybean [*Glycine max* (L.) Merr.] because, in part, of its weed-suppressive capabilities [93]. In Japan, a possible goal of a cover crop system in NT soybean is to eliminate or greatly reduce the use of herbicides in association with the implementation of reduced tillage and the appropriate use of cover crops [69]. Two annual medic species [burr medic (*M. polymorpha* cv. Santiago) and barrel medic (*M. truncatula* Gaertn. cv. Mogul)], berseem clover (*Trifolium alexandrinum* L. cv. Bigbee), and medium red clover (*Trifolium pratense* L.) were NT seeded as cover crops into winter wheat (*Triticum aestivum* L.) stubble in a winter wheat/corn (*Zea mays* L.) rotation system. The density of winter annual weeds was between 41 and 78 % lower following most cover crops when compared with no cover control in 2 out of 4 site years, while dry weight was between 26 and 80 % lower in all 4 trial site years [94].

In CA systems, the presence of crop residue acting as mulches on the soil surface, influence soil temperature and moisture regimes that affect weed

seed germination and emergence patterns over the growing season. Crop residues can influence weed seed germination and seedling emergence [95, 96]. Several mechanisms may contribute to reduced weed emergence and growth where surface cover crop residues are present, including reduction in light penetration to the soil [97], physical obstruction resulting in seed-reserve depletion before emergence [98], increased seed predation or decay [74, 99], decreased daily soil temperature fluctuations [101], or the production of allelopathic compounds [97].

Weed emergence generally declines with increasing residue amounts. However, the emergence of certain weed species is also favored by some crop residue at low amounts. For example, germination and growth of *Avena fatua* L. and *Avena sterilis* L. may get stimulated with low levels of wheat residue [11]. Further, late emerging weed plants produce fewer seeds than the early emerging ones [77]. For example, the residue of *Vicia villosa* Roth and *Secale cereal* L. reduced total weed density by more than 75 % compared with the treatments with no residue [100]. The presence of rye mulch in corn significantly reduced the emergence of *Chenopodium album* L., *Digitaria sanguinalis* (L.) Scop., and *Portulaca oleracea* L. and total weed biomass [101]. Tuesca *et al.* [102] attributed the lower densities of *Chenopodium album* L. under NT systems to the inhibitory effect of crop residues on light interception. Saini [103] found that rye residue provided 81–91% control of *Diodia virginiana* L. and *Jacquemontia tamnifolia* [L.] Griseb., *Digitaria sanguinalis* [L.] Scop. control was only 11 % in cotton and peanut.

For significant suppressive effects of mulch on the emergence and growth of *Echinochloa crusgalli* (L.) P. Beauv. and *Eclipta prostrata* (L.) L., 6 t ha⁻¹ rice residue as mulch was needed, whereas, the emergence of *Echinochloa colona* (L.) Link and *Dactyloctenium aegyptium* Willd. was reduced with as little as 1 to 2 t ha⁻¹ [68]. Chhokar *et al.* [104] observed that 2.5 t ha⁻¹ rice residue mulch was not effective in suppressing weeds, but 5.0 and 7.5 t ha⁻¹ residue mulch reduced weed biomass by 26 to 46 %, 17 to 55 %, 22 to 43 %, and 26 to 40 % of *Phalaris minor* Retz., *Oxalis corniculata*. L., *Medicago polymorpha* L. and *Setaria viridis* (L.) P.Beauv., respectively, compared with ZT without residue. Similar, Chauhan and Abugho [96] reported that 6 t ha⁻¹ crop residues reduced the emergence of *Echinochloa colona* (L.) Link, *Dactyloctenium aegyptium* Willd. and *Cyperus iria* L. by 80–95 %, but only reduce the emergence of *Echinochloa crusgalli* (L.) P. Beauv. by up to 35 %. However, crop residues alone may not be able to fully control

weeds, e.g. *Vicia villosa* residues suppressed weeds early in the growing season but herbicide was needed to achieve season-long weed control [11].

Unlike in the CT system, crop residues present at the time of herbicide application in CA systems may decrease the herbicide's effectiveness as the residues intercept the herbicide and reduce the amount of herbicide that can reach the soil surface and kill germinating seeds [51]. According to Chauhan *et al.* [105], crop residues can intercept 15–80 % of the applied herbicides and this may result in reduced efficacy of herbicides in CA systems.

DIVERSIFIED CROP ROTATION

Crop rotation increases crop diversity, reducing crop yield penalties associated with insect damage, diseases and weed infestations whilst improving nutrient cycling [38]. Crop rotations are arguably the most effective way to control weeds. It limits the build-up of weed populations and prevents weed shifts as the weed species tend to thrive in a crop with similar growth requirements. Different crops require different cultural practices, which help to disrupt the growing cycle of weeds and prevent any weed species to dominate [11]. In this way, any given crop can be thought of a filter, only allowing certain weeds to pass through its management regime [106]. Rotating crops will rotate selection pressures, preventing one weed from being repeatedly successful, and thus preventing its establishment [107]. Weed diversity has been shown to increase under crop rotation compared to monoculture [108, 109]. Greater diversity prevents the domination of a few problem weeds. Murphy *et al.* [110] observed the highest weed species diversity in NT fields with a three-crop rotation of corn–soybean–winter wheat. Weed species composition would be affected by rotation design, and weed population dynamics are very dependent on the crops included in the rotation [111]. The diversification of the system even for a short period and intensification by including summer legumes/green manuring decreased the weed menace [112]. The integration of red clover in the sweet corn–pea–wheat rotation led to a 96% reduction in the seed bank density of winter annuals [113]. Further, including perennial forages, such as alfalfa in rotation, has been shown to contribute weed control for up to three years, and can be particularly effective in NT systems [114, 115]. In NT systems of the Northern Great Plains of the United States (US) and Canada, stacked rotation designs offer superior weed control compared to yearly rotations [116]. Anderson and Beck [117] found that warm-season weeds were more prevalent in rota-

tions with two warm-season crops in 3 years, whereas these species were rare in rotations that included 2-year intervals of cool-season crops or fallow. Similar, weed community density declined across time with NT when rotations consisted of two cool-season crops followed by two warm-season crops; in contrast, weed community density was 13-fold greater with a two-crop rotation and NT [118]. In that context, in the semiarid Great Plains, producers who rotated cool-season and warm-season crops reduced weed community density and could grow some crops without needing herbicides to achieve optimum yields. With these diverse rotations, producers are using 50% less herbicide to manage weeds compared with that in less-diverse rotations [118].

HERBICIDES IN CA

Herbicides have an important role in weed control under CA systems [55]. Restricting tillage reduces weed control options and increases reliance on herbicides in such production systems, particularly the recent development of post-emergence broadspectrum herbicides provides an opportunity to control weeds in CA [119]. Therefore, CA is presently a common farming system in many countries, principally because many types of herbicides are available [120]. The presence of weed seeds on the upper soil surface, due to no-tillage operation, leads to higher weed infestation in CA, and so far herbicides are the only answer to deal with this problem [11]. They play an important role, particularly in controlling weeds during the first years after the adoption of conservation agriculture [121]. Herbicides are effective weed control measures and offer diverse benefits, such as saving labor and fuel cost, requiring less human efforts, reducing soil erosion, saving energy, increasing crop production, reducing the cost of farming, allowing flexibility in weed management, and tackling difficult-to-control weeds [122, 123]. The use of herbicides to facilitate weed control and soil cover management is an option to reduce production costs and to avoid the aforementioned negative effects associated with soil tillage, including the stimulation of further weed emergence and spread. In Canada adoption of NT has not increased herbicide use significantly [124], and in the US Great Plains, NT wheat systems have controlled weeds using cultural tactics and reduced herbicide usage by 50% compared to CT [118]. Similar, in some West European countries [125] and some areas in Australia [126], which have agroecological conditions similar to Europe, herbicide use

per tonne of output is lower in CA systems with integrated weed management than in CT farming [127]. Published research on NT corn and soybean showed that preemergence herbicide use for summer annual weed control could be reduced 50 % by banding herbicides over crop rows and substituting between row mowing for herbicides [128–130]. This equaled a 50 % reduction in preemergence herbicide use and a 25 % reduction in total herbicide use in no-till. In corn and soybean, crop yields were statistically indistinguishable among weed-free checks, broadcast preemergence herbicide treatments, and some treatments using banded preemergence herbicide followed by between-row mowing. Similarly, NT with effective herbicide weed control was more remunerative in the soybean–wheat system [131]. Further, the application of a burndown herbicide such as paraquat or glyphosate at planting followed by a herbicide such as Harmony Extra (Thifensulfuron-methyl+ Tribenuron-methyl) in the spring excellent solve the problems with broadleaved weeds no-till wheat [132]. Herbicide treatments such as glufosinate, mesotrione, and dicamba + diflufenzopyr are effective in suppressing *Taraxacum officinale* Web. competition in no-tillage corn [133].

However, to sustain CA systems, herbicide rotation and/or integration of weed management practices is preferred as continuous use of a single herbicide over a long period of time may result in the development of resistant biotypes, shifts in weed flora and negative effects on the succeeding crop and environment [11, 134]. Selection pressure imparted by herbicide tactics can result in weed shifts attributable to the natural resistance of a particular species to the herbicide or the evolution of herbicide resistance within the weed population [55]. Reddy [135] found that continuous bromoxynil-resistant cotton production resulted in weed species shift toward *Portulaca oleracea* L., *Senna obtusifolia* [L.] Irwin and Barneby, and (*Cyperus esculentus* L.).

A major criticism of CA is its enhanced reliance on herbicides as compared to tilled systems. In particular, glyphosate may be heavily used, especially to control perennial weeds [136]. When herbicides are not used at appropriate rates or in rotation [40], it can lead to the development of herbicide resistance among major weed species [137,138]. Herbicide resistance and weed control problems are clearly the main reasons given by adopters for past or intended reductions in NT use [139]. Herbicide resistance and weed control issues are the major reason why some NT adopters are reducing their use of NT as the core principle of CA.

REFERENCES

- [1] D. R. Shaw, S. Culpepper, M. Owen, A. Price, R. Wilson, Herbicide-resistant weeds threaten soil conservation gains: Finding a balance for soil and farm sustainability. Council for agricultural science and technology, Issue paper 49, Ames, Iowa, 2012.
- [2] M. A. Sprague, Overview, In: M. A. Sprague and G. B. Triplett (eds.). *No-tillage and surface-tillage agriculture*, Wiley, New York, 1986, pp. 1–18.
- [3] D. C. Reicosky, R. R. Allmaras, Advances in tillage research in North American cropping systems, *J. Crop Prod.*, **8** (2003), pp.75–125.
- [4] R. J. Godwin, Potential of “No-till” systems for arable farming. The Worshipful company of farmers soil and water management centre Harper Adams, University Newport Shropshire TF108NB, 2014. <https://cdn.harper-adams.ac.uk/document/project/Potential-of-No-till-Systems-for-Arable-Farming-Report.pdf>
- [5] P. C. Wall, Tailoring conservation agriculture to the needs of small farmers in developing countries: An analysis of issues. *J. Crop Improve.*, **19** (2007), pp.137–155.
- [6] C. Thierfelder, P. C. Wall, Effects of conservation agriculture techniques on infiltration and soil water content in Zambia and Zimbabwe. *Soil Till. Res.*, **105** (2009), pp.217–227.
- [7] M. U. F. Kirschbaum, The temperature dependence of soil organic matter decomposition, and the effect of global warming on soil organic C storage. *Soil Biol. Biochem.*, **27** (1995), pp. 753–760.
- [8] S. J. Del Grosso, A. R. Mosier, W. J. Parton, D. S. Ojima, DAYCENT model analysis of past and contemporary soil NO and net greenhouse gas flux for major crops in the USA. *Soil and Tillage Research*, **83** (2005), pp. 9–24.
- [9] C. Johansen, M. E. Haque, R. W. Bell, C. Thierfelder, R. J. Esdaile, Conservation agriculture for small holder rainfed farming: opportunities and constraints of new mechanized seeding systems. *Field Crops Res.*, **132** (2012), pp. 18–32.
- [10] C. Thierfelder, P. C. Wall, Effects of conservation agriculture on soil quality and productivity in contrasting agro-ecological environments of Zimbabwe. *Soil Use Manag.*, **28** (2012), pp. 209–220.
- [11] V. Singh, K. Barman, R. Singh, A. Sharma, Weed management in conservation agriculture systems, In: Farooq M., Siddique K. (eds) *Conservation Agriculture*, Springer, Cham, 2015.
- [12] R. Derpsch, History of crop production with and without tillage, leading edge, *The Journal of No-Till Agriculture*, **3** (2004), pp. 150-154.
- [13] O. C. Burnside, The history of 2,4-D and its impact on development of the discipline of weed science in the United States, In O. C. Burnside, ed. *Biologic and Economic Assessment of Benefits from Use of Phenoxy Herbicides in the United States*, Washington, DC: U.S. Department of Agriculture NAPIAP Rep. 1-PA-96, 1996, p. 5–15.
- [14] R. Derpsch, T. Friedrich, A. Kassam, L. Hongwen, Current status of adoption of no-till farming in the world and some of its main benefits, *Int. J. Agric. Biol.Eng.*, **3** (2010), pp. 1–25.
- [15] A. R. Sharma, V. P. Singh, Integrated weed management in conservation agriculture systems, *Indian Journal of Weed Science*, **46** (2014) pp. 23–30.
- [16] FAO (Food and Agriculture Organization), Introduction to conservation agriculture (its principles and benefits), Rome, March, 2013. <http://teca.fao.org/technology/introduction-conservationagriculture-its-principles-benefits>
- [17] R. Derpsch, The extent of conservation agriculture adoption worldwide: implications and impact, *Proceedings 3rd World Congress on Conservation Agriculture*, Nairobi, 2005, CDrom, p. 15.
- [18] P. R. Hobbs, Conservation agriculture: What is it and why is it important for future sustainable food production? *J. Agric. Sci.*, **145** (2007) pp.127–137.
- [19] T. Friedrich, R. Derpsch, A. Kassam, Overview of the global spread of conservation agriculture. Special Issue 6, reconciling poverty eradication and protection of the environment. *Field Actions Science Reports*, 2012.
- [20] Anonymous, Conservation agriculture. Division of Statistics and Strategic Foresight – Centre for Studies and Strategic Foresight Analysis No. 61, 2013. www.agreste.agriculture.gouv.fr
- [21] P. R. Hobbs, K. Sayre, R. Gupta, The role of conservation agriculture in sustainable agriculture. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, **363** (2008), pp. 543–555.
- [22] T. Friedrich, A. H. Kassam, Adoption of conservation agriculture technologies: constraints and opportunities, Invited paper, *IV World Congress on Conservation Agriculture*, New Delhi, India, 2009a, 4-7 February.
- [23] T. Friedrich, A. H. Kassam, F. Taher, Adoption of conservation agriculture and the role of policy and institutional support, Invited keynote paper presented at the *International Consultation on No-Till with Soil Cover and Crop Rotation: A Basis for Policy Support to Conservation Agriculture for Sustainable Production Intensification*, Astana-Shortandy, Kazakhstan, 2009b, July.
- [24] T. Friedrich, J. Kienzle, A. H. Kassam, Conservation agriculture in developing countries: The role of mechanization, Paper presented at the *Club of Bologna Meeting on Innovation for Sustainable Mechanisation*, Hanover, Germany, 2009c, 2nd November.
- [25] M. Lane, M. Kibblewhite, L. Montanarella, Conservation agriculture in Europe. An approach to

- sustainable crop production by protecting soil and water? SOWAP, Jealott's Hill International Research Centre, Bracknell, Berkshire RG42 6EY, 2006.
- [26] N. Verhulst, B. Govaerts, E. Verachtert, A. Castellanos-Navarrete, M. Mezzalama, P. Wall, J. Deckers, K.D. Sayre, Conservation agriculture, improving soil quality for sustainable production systems? In: Lal, R., Stewart, B.A. (Eds.), *Advances in Soil Science: Food Security and Soil Quality*, CRC Press, Boca Raton, FL, USA, 2010, p. 137–208.
- [27] FAO, (Food and Agriculture Organization), Save and Grow: A policymaker's guide to the sustainable intensification of small holder crop production, Food and Agriculture Organization of the United Nations, Rome, 2011.
- [28] J. L. Steiner, H.H. Schomberg, P.W. Unger, J. Cresap, Biomass and residue cover relationships of fresh and decomposing small grain residue. *Soil Science Society of America Journal*, **64** (2000), pp. 2109–2114.
- [29] O. Erenstein, K. Sayre, P. Wall, J. Hellin, J. Dixon, Conservation agriculture in maize- and wheat-based systems in the (sub)tropics: lessons from adaptation initiatives in South Asia, Mexico, and Southern Africa. *J. Sustain.Agric.*, **36** (2012), pp. 180–206.
- [30] L. García-Torres, A. Martínez-Vilela, F.S. De Noreña conservation agriculture in Europe: Current status and perspectives. In: García-Torres L., Benites J., Martínez-Vilela A., Holgado-Cabrera A. (eds) *Conservation Agriculture*, Springer, Dordrecht, 2003.
- [31] E. J. González-Sánchez, R. Ordoñez-Fernández, R. Carbonell-Bojollo, O. Veroz-Gonzalez, J.A. Gil-Ribes, Meta-analysis on atmospheric carbon capture in Spain through the use of conservation agriculture. *Soil Tillage*, **122** (2012), pp. 52–60.
- [32] Y. Gao, X.H. Dang, Y. Yu, Effects of tillage methods on soil carbon and wind erosion. *Land Degrad. Dev.*, **27** (2016), pp. 583–591.
- [33] J. Six, S. M. Ogle, F. J. Breidt, R. T. Conant, A. R. Mosier, K. Paustian, The potential to mitigate global warming with no-tillage is only realized when practiced in the long-term. *Glob. Change Biol.*, **10** (2004), pp. 155–160.
- [34] V. Chaplota, C. N. Mchunub, A. Mansonc, S. Lorentz, G. P. W. Jewitt, Water erosion-induced CO₂ emissions from tilled and no-tilled soils and Sediments. *Agric Ecosyst Environ.*, **159** (2012), pp. 62–69.
- [35] C. Van Kessel, R. Venterea, J. Six, M. Adviento-Borbe, B. A. Linnquist, K. J. Van Groenigen, Climate duration and N placement determine N₂O emissions in reduced tillage systems: a meta-analysis. *Glob. Change Biol.*, **19** (2013), pp. 33–44.
- [36] R. Lal, Enhancing crop yields in the developing countries through restoration of the soil organic carbon pool in agricultural lands. *Land Degrad. Dev.* **17** (2005), pp. 197–209.
- [37] B. Govaerts, M. Mezzalama, Y. Unno, S. D. Sayre, M. Luna-Guido, K. Vanherck, J. Deckers, Influence of tillage, residue management, and crop rotation on soil microbial biomass and catabolic diversity. *Appl. Soil Ecol.*, **37**, (2007), pp. 18–30.
- [38] R. Derpsch, T. Friedrich, Global overview of conservation agriculture adoption. In *Proceedings of the 4th World Congress on Conservation Agriculture*, New Delhi, India, 4–7 February 2009a, pp. 429–438.
- [39] L. C. Standifer, C. E. Beste, Weed control methods for vegetable production with limited-tillage, In: A. F. Wiese. (ed.). *Weed control in limited tillage systems*, *Weed Sci. Soc. Amer. Monogr.*, **2** (1985), pp. 93–100.
- [40] S. B. Powles, Q. Yu, Evolution in action: plants resistant to herbicides. *Annu. Rev. Plant. Biol.*, **61**, (2010), 317–347.
- [41] G. D. Hoyt, T. R. Konsler, Soil water and temperature regimes under tillage and cover crop management for vegetable culture. *Proc. 11th Intl. Conf., ISTRO*, Edinburgh, Scotland, 1988, p. 697–702.
- [42] B. D. Soane, B. C. Ball, J. Arvidsson, G. Basch, F. Moreno, J. Roger- Estrande, No-till in northern, western and south-western Europe: A review of problems and opportunities for crop production and the environment. *Soil Tillage and Research*, **118** (2012), pp. 66–87.
- [43] S. H. Phillips, Introduction in No-tillage agriculture: Principles and practice. Van Nostrand Reinhold, New York, 1984, p. 1–10.
- [44] D. R. Huggins, J.P. Reganold, No-till: the quiet revolution. *Scientific American*, July 2008, pp. 70–77.
- [45] A. R. Sharma, M. L. Jat, Y. S. Saharawat, V. P. Singh, R. Singh, Conservation agriculture for improving productivity and resource-use efficiency: Prospects and research needs in Indian context. *Indian Journal of Agronomy*, **57**(3rd IAC Special Issue): (2012), pp. 131–140.
- [46] C. M. Pittelkow, X. Liang, B. A. Linnquist, K. J. Van Groenigen, J. Lee, M. E. Lundy, N. van Gestel, J. Six, R. T. Venterea, C. van Kessel, Productivity limits and potentials of the principles of conservation agriculture, *Nature*, **517**, (2015), pp. 365–368.
- [47] D. A. Derksen, R. L. Anderson, R. E. Blackshaw, B. Maxwell, Weed dynamics and management strategies for cropping systems in the northern Great Plains, *Agron. J.*, **94** (2002), pp. 174–185.
- [48] K. E. Giller, E. Witter, M. Corbeels, P. Tittonell, Conservation agriculture and small holder farming in Africa: the heretics' view. *Field Crops Res.*, **114**, (2009), pp. 23–34.
- [49] M. Farooq, K. Flower, K. Jabran, A. Wahid, K. H. Siddique, Crop yield and weed management in

- rained conservation agriculture. *Soil Tillage Res.* **117** (2011), pp.172–183.
- [50] L. O. Pedro, A. Machado de, C. A. Silva, Soil management under no-tillage systems in the tropics with special reference to Brazil, *Nutrient Cycling in Agroecosystems*, **61** (2001), pp.119–130.
- [51] A. Shrestha, T. Lanini, S. Wright, R. Vargas, J. Mitchell, Conservation Tillage and Weed Management. *ANR Publication 8200.Regents of the University of California Division of Agriculture and Natural Resources*, 2006. <https://anrcatalog.ucanr.edu/pdf/8200.pdf>
- [52] D. R. Clements, D. L. Benoit, S. D. Murphy, C. J. Swanton, Tillage effects on weed seed return and seed bank composition. *Weed Sci.*, **44** (1996), pp. 314–322.
- [53] R. G. Hartzler, M. D. Owen, Weed management in conservation tillage systems. Ames: Iowa State University, University Extension, 1997. <http://www.extension.iastate.edu/Publications/PM1176.pdf>
- [54] C. J. Swanton, A. Shrestha, S. Z. Knezevic, R. C. Roy, B.R. Ball-Coelho, Influence of tillage type on vertical weed seed bank distribution in a sandy soil. *Can. J. Plant Sci.*, **80** (2000), pp. 455–457.
- [55] S. V. Eslami, Weed management in conservation agriculture systems, Chapter 5, In book: *Recent Advances in Weed Management*, 2014.
- [56] J. S. Mishra, V. P. Singh Tillage and weed control effects on productivity of a dry seeded rice–wheat system on a vertisol in central India. *Soil and Tillage Research*, **123** (2012), pp. 11–20.
- [57] E. Demjanová, M. Macák, I. Đalović, F. Majerník, Š. Týr, J. Smatana, Effects of tillage systems and crop rotation on weed density, weed species composition and weed biomass in maize, *Agronomy Research*, **7** (2009), pp. 785–792.
- [58] M. Tolimir, M. Vesković, I. Komljenović, I. Djalović, B. Stipešević, Influences of soil tillage and fertilization on maize yield and weed infestation, *Cereal Res. Comm.*, **34**, (2006), pp. 323–326 part I.
- [59] C. J. Swanton, A. Shrestha, S. Z. Knezevic, R. C. Roy, B. R. Ball-Coelho, Effect of tillage systems, N, and cover crop on the composition of weed flora. *Weed Sci.*, **47** (1999), pp. 454–461.
- [60] D. A. Derksen, G. P. Lafond, A. G. Thomas, H. A. Loepky, C. J. Swanton, Impact of agronomic practices on weed communities: Tillage systems. *Weed Sci.*, **41** (1993), pp. 409–417.
- [61] R. J. Froud-Williams, Changes in weed flora with different tillage and agronomic management systems. In M. A. Altieri and M. Liebman, eds., *Weed Management in Agroecosystems: Ecological Approaches*. Boca Raton: CRC Press, 1988, p. 213–236.
- [62] D. Childs, T. Jordan, M. Ross, T. Bauman, Weed control in no-tillage systems. Purdue University Cooperative Extension Service. Conservation Tillage Series CT-2. West Lafayette: Purdue University, 2001.
- [63] J. Cardina, C. P. Herms, D. J. Doohan, Crop rotation and tillage system effects on weed seedbanks. *Weed Sci.*, **50** (2002), pp. 448–460.
- [64] W. S. Curran, D. D. Lingenfelter, L. Garling, An introduction to weed management for conservation tillage systems. Conservation Tillage Series No. 2. College of Agricultural Sciences, Cooperative Extension. University Park, PA: The Pennsylvania State University, 1996.
- [65] Y. Gan, N. K. Harker, B. McConkey, M. Suleimenov, Moving towards no-till practices in Northern Eurasia. In: *Goddard, T., Zoebisch, M., Gan, Y., Ellis, W., Watson, A., Sombatpanit, S. (Eds.), No-till Farming Systems (Paraguay)*, 2008.
- [66] B. Frick, A. G. Thomas, Weed surveys in different tillage systems in southwestern Ontario field crops. *Can J Plant Sci.*, **72** (1992), pp. 1337–1347.
- [67] V. Kumar, J. K. Ladha, Direct-seeding of rice: recent developments and future research needs. *Adv. Agron.* **111** (2011), pp.297–413.
- [68] V. Kumar, S. Singh, R. S. Chhokar, R. K. Malik, D. C. Brainard, J. K. Ladha, Weed management strategies to reduce herbicide use in zero-till rice–wheat cropping systems of the Indo-Gangetic plains. *Weed Technology*, **27** (2013), pp. 241–254.
- [69] H. Kobayashi, S. Miura, A. Oyanagi, Effects of winter barley as a cover crop on the weed vegetation in a no-tillage soybean, *Weed Biology and Management*, **4** (2004), pp. 195–205.
- [70] D. A. Ball, Weed seed bank response to tillage, herbicide, and crop rotation sequence, *Weed Sci.*, **40** (1992), pp. 654–659.
- [71] M. R. Pareja, D.W. Staniforth, Distribution of weed seed among soil structural units, *Weed Sci.*, **33** (1985), pp.182–189.
- [72] O. Auškalnienė, A. Auškalnis, The influence of tillage system on diversities of soil weed seed bank, *Agronomy Research*, **7** (special iss. 1) (2009), pp.156–161.
- [73] B. S. Chauhan, G. S. Gill, C. Preston, Tillage systems affect trifluralin bioavailability in soil, *Weed Sci.*, **54** (2006), pp. 941–947.
- [74] H. E. Cromar, S. D. Murphy, C. J. Swanton Influence of tillage and crop residue on post dispersal predation of weed seeds, *Weed Sci.*, **47** (1999), pp. 184–194.
- [75] A. Davis, K. Renner, C. Sprague, L. Dyer, D. Mutch, Integrated weed management: one year’s seeding... East Lansing: Michigan State University. Extension Bulletin E-2931, 2005.
- [76] H. S. Jacob, D. M. Minkey, R. S. Gallagher, C. P. Borger, Variation in post dispersal weed seed pre-

- dition in a crop field. *Weed Sci.*, **54** (2006), pp.148–155.
- [77] B. S. Chauhan, D. E. Johnson, Implications of narrow crop row spacing and delayed *Echinochloa colona* and *Echinochloa crus-galli* emergence for weed growth and crop yield loss in aerobic rice, *Field Crops Research*, **117** (2010), pp. 177–182.
- [78] C. Mwale, Effect of Tillage Practices on Weed Populations and Seed Banks in Maize based Production Systems in Malawi (Master thesis), University of Lyon, ISARA-Lyon, (2009), pp. 1–45.
- [79] G. R. Sagar, A. M. Mortimer, An approach to the study of the population dynamics of plants with special reference to weeds, *Adv. Appl. Biol.*, **1** (1976), pp.1–47.
- [80] K. Usman, S. K. Khalil, M. A. Khan, Impact of tillage and herbicides on weed density and some physiological traits of wheat under rice-wheat cropping system. *Sarhad J. Agric.*, **26** (2010), pp. 475–487.
- [81] R. E. Peachey, C. Mallory-Smith, Influence of winter seed position and recovery date on hairy nightshade (*Solanum sarrachoides*) recruitment and seed germination, dormancy, and mortality, *Weed Sci.*, **55** (2007), pp. 49–59.
- [82] R. E. Peachey, C. Mallory-Smith Effect of fall tillage and cover crop strategies on wild proso millet (*Panicum miliaceum*) emergence and interference in snap beans. *Weed Technol.*, **25** (2011), pp. 119–126.
- [83] D. R. Clements, S. F. Weise, C. J. Swanton, Integrated weed management and weed species diversity, *Phytoprotection*, **75** (1994), pp. 1–18.
- [84] R. Skuodienė, D. Karčauskienė, S. Čiuberkis, R. Repšienė, D. Ambrazaitienė, The influence of primary soil tillage on soil weed seed bank and weed incidence in a cereal-grass crop rotation, *Zemdirbyste-Agriculture*, **100** (2013), pp. 25–32.
- [85] F. D. Menalled, K. L. Gross, M. Hammond, Weed above ground and seed bank community responses to agricultural management systems, *Ecological Applications*, **11** (2001), pp. 1586–1601.
- [86] A. Isensee, A. Sadeghi, Effects of tillage and rainfall on atrazine residue levels in soil, *Weed Science*, **42** (1994), pp. 462–467.
- [87] G. G. Nagabhushana, A. D. Worsham, J. P. Yenish. Allelopathic cover crops to reduce herbicide use in sustainable agriculture systems. *Allelopathy J.*, **8** (2001), pp.133–146.
- [88] S. C. Phatak, Managing pests with cover crops, In C. Shirley, G. Bowman, and C. Cramer, eds. *Managing Cover Crops Profitably*, Burlington, VT: *Sustainable Agriculture*, (1998), pp. 25–33.
- [89] M. E. Foley, Genetic approach to the development of cover crops for weed Management, *Journal of Crop Production*, **2** (1999), pp. 77–93.
- [90] A. J. Price, M. E. Stoll, J. S. Bergtold, F. J. Arriaga, K. S. Balkcom, T.S. Kornecki, R.L. Raper, Effect of cover crop extracts on cotton and radish radical elongation, *Communications in Biometry and Crop Science*, **3** (2008), pp. 60–66.
- [91] M. C. Akemo, E. E. Regnier, M. A. Bennett, Weed suppression in spring-sown rye (*Secale cereale*)–pea (*Pisum sativum*) cover crop mixes, *Weed Technol.*, **14** (2000), pp.545–549.
- [92] J. P. Yenish, A. D. Worsham, A. C. York, Cover crops for herbicide replacement in no-tillage corn (*Zea mays*), *Weed Technol.*, **10** (1996) pp.815–821.
- [93] R. Derpsch, C. H. Roth, N. Sidiras, U. Kopke, Controle da erosao no Parana', Brazil: Sistemas de cobertura do solo, plantio directo e prepare conservacionista do solo. GmbH, Eschborn, Germany: Deutsche Gesellschaft fur Technische Zusammenarbeit (GTZ), 1991.
- [94] J. W. Fisk, O. B. Hesterman, A. Shrestha, J. J. Kells, R. R. Harwood, J. M. Squire, C. C. Sheaffer, Weed suppression by annual legume cover crops in no-tillage corn, *Agronomy Journal*, **93** (2000), pp. 319–325.
- [95] B. S. Chauhan, Weed ecology and weed management strategies for dry-seeded rice in Asia, *Weed Technol.*, **26** (2012), pp.1–13.
- [96] B. S. Chauhan S. B. Abugho, Interaction of rice residue and PRE herbicides on emergence and biomass of four weed species, *Weed Technol.*, **26** (2012), pp. 627–632.
- [97] J. R. Teasdale, C. L. Mohler, Light transmittance, soil temperature and soil moisture under residue of hairy vetch and rye, *Agron J.*, **85** (1993), pp. 673–680.
- [98] J. R. Teasdale, C. L. Mohler, The quantitative relationship between weed emergence and the physical properties of mulches, *Weed Sci.*, **48** (2000), pp. 385–392.
- [99] D. C. Brainard, R. E. Peachey, E. R. Haramoto, J.M. Luna, A. Rangarajan Weed ecology and nonchemical management under strip-tillage: implications for northern U.S. vegetable cropping systems, *Weed Technol.*, **27** (2013), pp. 218–230.
- [100] C. L. Mohler, J. R. Teasdale, Response of weed emergence to rate of *Vicia villosa* Roth and *Secale cereale* L. residue, *Weed Research*, **33** (1993), pp. 487–499.
- [101] C. L. Mohler, M. B. Callaway, Effects of tillage and mulch on the emergence and survival of weeds in sweet corn, *Journal of Applied Ecology*, **29** (1992), pp. 21–34.
- [102] D. Tuesca, E. Puricelli, J. C. Papa, A long-term study of weed flora shifts in different tillage systems, *Weed Res.*, **41** (2001), pp. 369–382.
- [103] M. Saini, Integration of cover crop residues, conservation tillage and herbicides for weed manage-

- ment in corn, cotton, peanut and tomato, Dissertation, Auburn University, Auburn, Alabama, 2009.
- [104] R. S. Chhokar, S. Singh, R. K. Sharma, M. Singh, Influence of straw management on *Phalaris minor* control, *Indian Journal of Weed Science*, **41** (2009), pp. 150–156.
- [105] B. S. Chauhan, R. G. Singh, G. Mahajan, Ecology and management of weeds under conservation agriculture: A review. *Crop Protection*, **38** (2012), pp. 57–65.
- [106] B. D. Booth, C. J. Swanton, Assembly theory applied to weed communities *Weed Sci.*, **50** (2002), pp. 2–13.
- [107] V. Nichols, N. Verhulst, R. Cox, B. Govaerts, Weed dynamics and conservation agriculture principles: A review, *Field Crops Research*, **183** (2015), pp. 56–68.
- [108] F. C. Stevenson, A. Légère, R. R. Simard, D. A. Angers, D. Pangeau, J. Lafond, Weed species diversity in spring barley varies with crop rotation and tillage, but not with nutrient source, *Weed Sci.*, **45** (1997), pp. 798–806.
- [109] D. L. Karlen, G. E. Varvel, D. G. Bullock, R. M. Cruse, Crop rotations for the 21st century. *Adv Agron.*, **53** (1994), pp. 1–45.
- [110] S. D. Murphy, D. R. Clements, S. Belaoussoff, P. G. Kevan, C. J. Swanton, Promotion of weed species diversity and reduction of weed seedbanks with conservation tillage and crop rotation, *Weed Sci.*, **54** (2006), pp. 69–77.
- [111] J. Dorado, J. P. del Monte, C. López-Fando, Weed seed bank response to crop rotation and tillage in semiarid agroecosystems, *Weed Sci.*, **47** (1999), pp. 67–73.
- [112] R. K. Singh, J. S. Bohra, V. K. Srivastava, R. P. Singh, Effect of diversification of rice–wheat system on weed dynamics in Rice, *Indian Journal of Weed Science*, **40** (2008), pp. 128–131.
- [113] D. C. Brainard, R. R. Bellinde, R. R. Hahn, D. A. Shah, Crop rotation, cover crop, and weed management effects on weed seed banks and yields in snap bean, sweet corn, and cabbage, *Weed Science*, **56** (2008), pp. 434–441.
- [114] P. Ominski, M. Entz, N. Kenkel, Weed suppression by *Medicago sativa* in subsequent cereal crops: a comparative survey, *Weed Sci.*, **47** (1999), pp. 282–290.
- [115] P. Ominski, M. Entz, Eliminating soil disturbance reduces post-alfalfa summer annual weed populations, *Can. J. Plant Sci.*, **81** (2001), pp. 881–884.
- [116] R. L. Anderson, Sequencing crops to minimize selection pressure for weeds in the central great plains, *Weed Technol.*, **18** (2004), pp. 157–164.
- [117] R. L. Anderson D. L. Beck, Characterizing weed communities among various rotations in central South Dakota, *Weed Technol.*, **21** (2007), pp. 76–79.
- [118] R. L. Anderson, A multi-tactic approach to manage weed populations in crop rotations, *Agron. J.*, **97** (2005), pp. 1579–1583.
- [119] G. Mahajan, L. S. Brar, V. Sardana, Effect of tillage and time of sowing on the efficacy of herbicides against *Phalaris minor* in wheat, (In): *Proceeding of 17th Asian Pacific Weed Science Society Conference*, Bangkok, Thailand, (1999), pp. 193–198.
- [120] W. C. Koskinen, C. G. McWhorter Weed control in conservation tillage, *J. Soil Water Cons.*, **41** (1986), pp. 365–370.
- [121] M. M. Rahman, Weed management in conservation agriculture, *Adv Plants Agric Res.*, **7** (2017) 00253.
- [122] Z. Pacanoski, Herbicide use: benefits for society as a whole- a review, *Pak J Weed Sci Res.*, **13** (2007), pp. 135–147.
- [123] Z. Pacanoski, Neglected side of the coin. The real image of herbicides, Lambert Academic Publishing, Germany, 2018.
- [124] D. Derksen, R. Blackshaw, S. Boyetchko, Sustainability, conservation tillage and weeds in Canada, *Can. J. Plant Sci.*, **76** (1996), pp. 651–659.
- [125] V. Brautigam, F. Tebrugge, Influence of long-termed no-tillage on soil borne plant pathogens and on weeds, In: Tebrugge, F., Bohrsen, A. (Eds.), Experiences with the Applicability of No-tillage Crop Production in the West-European Countries, *Proc. EC Workshop III*, Wissenschaftlicher Fachverlag, Giessen, Germany, 1997, p. 17–29.
- [126] B. Crabtree, Weed control is superior with no-till. In: Crabtree, B. (ed.) Search for Sustainability with No-Till Bill in Dryland Agriculture. Beckenham, W.A.: Crabtree Agricultural Consulting, 2010, pp. 27–36.
- [127] J. F. C. Barros, G. Basch, M. Carvalho, Effect of reduced doses of a postemergence graminicide to control *Avena sterilis* L. and *Lolium rigidum* G. in no-till wheat under Mediterranean environment. *Crop Prot.*, **27** (2008), pp. 1031–1037.
- [128] W. W. Donald, Between row mowing + in-row band-applied herbicide for weed control in *Glycine max*, *Weed Sci.*, **48** (2000 a), pp. 487–500.
- [129] W. W. Donald, Timing and frequency of between-row mowing and band-applied herbicide for annual weed control in soybean, *Agron. J.*, **92** (2000 b), pp. 1013–1019.
- [130] W. W. Donald, N. R. Kitchen, K. A. Sudduth, Between-row mowing + banded herbicide to control annual weeds and reduce herbicide use in no-till soybean (*Glycine max*) and corn (*Zea mays*). *Weed Technol.*, **15** (2001), pp. 576–584.
- [131] J. S. Mishra, V. P. Singh, Weed dynamics and productivity of soybean (*Glycine max*) - based cropping systems as influenced by tillage and weed management. *Indian Journal of Agronomy*, **54** (2009), pp. 29–35.

- [132] J. Herbek, L. Murdock, J. Grove, L. Grabau, D. Van Sanford, J. Martin, J. James, D. Call, Comparing No-Till and Tilled Wheat in Kentucky. Cooperative Extension Service, University of Kentucky College of Agriculture, Lexington, KY, 40546, 2009.
- [133] A. S. Franssen, J. J. Kells, Common dandelion (*Taraxacum officinale*) control with postemergence herbicides in no-tillage glufosinate-resistant corn, *Weed Technology*, **21** (2007), pp. 14–17.
- [134] G. B. Triplett Jr, W. A. Dick No-tillage crop production: a revolution in agriculture. *Agron J.*, **100** (Suppl) (2008) S-153-S-115.
- [135] K.N. Reddy, Weed control and species shift in bromoxynil- and glyphosate-resistant cotton (*Gossypium hirsutum*) rotation systems, *Weed Technol.*, **18** (2004), pp. 131–139.
- [136] J. Moyer, E. Roman, C. Lindwall, R. Blackshaw, Weed management in conservation tillage systems for wheat production in North and South America, *Crop Prot.*, **13** (1994), pp. 243–259.
- [137] M. J. Owen, M. J. Walsh, R. S. Llewellyn, S. B. Powles, Widespread occurrence of multiple herbicide resistance in Western Australian annual ryegrass (*Lolium rigidum*) populations. *Crop Pasture Sci.*, **58** (2007), pp.711–718.
- [138] S. B. Powles, Evolved glyphosate-resistant weeds around the world: lessonst o be learned, *Pest Manag. Sci.*, **64** (2008), pp. 360–365.
- [139] F. H. D’Emden, R. S. Llewellyn, No-tillage adoption decisions in southern Australian cropping and the role of weed management. *Australian Journal of Experimental Agriculture*, **46** (2006), pp. 563–569.

ВЛИЈАНИЕ НА ТРИТЕ СТОЛБА НА КОНЗЕРВАЦИСКОТО ЗЕМЈОДЕЛСТВО И ХЕРБИЦИДИТЕ ВРЗ ПЛЕВЕЛИТЕ

Звонко Пацаноски

Факултет за земјоделски науки и храна, Универзитет „Св. Кирил и Методиј“,
Скопје, Република Македонија

Конзервациското земјоделство е засновано на три големи столба: редуцирана обработка или целосно нејзино елиминирање, покриеност на почвата и разновиден плодоред. Еден од најголемите проблеми со кои се соочува овој систем се плевелите и правилната борба против нив. Конзервацискиот систем предизвикува промена во динамиката на плевелните семиња во почвата, плевелната популација, заплевеленоста и растот на плевелите, со што се наметнува потребата од усогласување и изработување стратегија за борба против нив. Растителните остатоци влијаат врз ртењето на плевелите и нивните поници преку попречување на достапноста на светлината, физичкото задушвање и хемиското (алелопатско) дејствување. Покривните култури се основен и одржлив начин за физичко и хемиско (алелопатско) намалување на бројноста на плевелите. Разновидноста на културите во плодоредот помага во прекинување на циклусот на пораст на плевелите и ја спречува појавата на доминација на некои плевелни видови. Хербицидите, исто така, претставуваат интегрален дел од борбата против плевелите во конзервациското земјоделско производство. Но, кога не се употребуваат во препорачаните дози или во хербициден плодоред, можат да предизвикаат загадување на животната средина, промени во плевелната популација и појава на резистентност кај некои плевелни видови.

Клучни зборови: конзервациско земјоделско производство; плевели; растителни остатоци; покривни култури; плодоред