



Revista
Stomatologjike
Shqiptare

Revista Stomatologjike Shqiptare

Periodik
i Fakultetit të Mjekësisë Dentare



Revista Stomatologjike Shqiptare është revista zyrtare e Fakultetit të Mjekësisë Dentare, Universiteti i Mjekësisë, Tiranë. Revista botohet në gjuhën shqipe dhe angleze, dy herë në vit.

Revista Stomatologjike Shqiptare ka për qëllim të botojë materiale me cilësi të lartë shkencore dhe klinike në të gjitha fushat e Stomatologjisë. Materialet e paraqitura për botim shqyrtohen nga redaksia e Revistës. Ajo synon gjithashtu të informojë mjekët stomatologë mbi aktivitete dhe të reja, si dhe të ofrojë një mundësi debati mbi aspekte të ndryshme të stomatologjisë.

The Albanian Dental Journal is the official Journal of the Dental Medicine Faculty at the Medical University of Tirana, Albania, published twice a year in Albanian and English language.

It is a peer-reviewed journal aiming to publish high quality material, both clinical and scientific, on all aspects of Dentistry. In addition it provides a mean of information for dentists about news and events and also a forum for exchange of opinions on all aspects of dentistry, including education issues.

Bordi Editorial Kombëtar / National Editorial Board

| | |
|------------------|------------------|
| Xhina Mulo | Rozela Xhemnica |
| Ramazan Isufi | Enida Petro |
| Lindita Xhemnica | Fatmir Lela |
| Adem Alushi | Kreshnik Keraj |
| Rozarka Budina | Silvana Bara |
| Ruzhdie Qafmolla | Esat Bardhoshi |
| Fejzi Keraj | Prunela Poliçi |
| Diana Brovina | Andis Qendro |
| Edit Xhajanka | Almira Isufi |
| Besnik Gavazi | Manola Kelmendi |
| Merita Bardhoshi | Vladimir Filaj |
| Ilma Robo | Matilda Imeraj |
| Nikollë Deda | Neada Hysenaj |
| Koço Gjilo | Anis Thodhorjani |
| Alketa Qafmolla | |

Bordi Editorial Ndërkombëtar / International Editorial Board

| | |
|---------------------------|---------------------|
| Sergio Caputi | Gianna Di Palma |
| Rosa Chaviano Moran | Giuseppina Malcangi |
| Christof Pertl | Dubravka Knezovic |
| Florian Beuer | Anna Minovska |
| Peter Pospiech | Bruna Sinjari |
| Norina Forna | Vincenzo Campanella |
| Giovanni Arcudi | Gianmaria D'Addazio |
| Francesco Inchingolo | Kenan Ferati |
| Giovanni Arcuri | Arbëresha Ferati |
| Giulia Donadel | Teuta Pustina |
| Lorenzo Arcuri | Fatmir Dragidella |
| Marco Jakovac | Manlio Santilli |
| Angelo Inchingolo | Imena Rexhepi |
| Giovanni Manes Gravina | |
| Alessio Danilo Inchingolo | |

Zyra e redaksisë

Editorial Office

Genc Kruja

Marsela Fagu

Lorena Qafmolla

Adresa: Fakulteti i Mjekësisë Dentare, Universiteti i Mjekësisë, Rr. Dibrës, 371, Tiranë, Shqipëri.

Address: Faculty of Dental Medicine at the University of Medicine, Tirana, Rr. Dibrës, 371, Tirana, Albania.

e-mail: redaksia_rssh@yahoo.com



Përmbajtja/Table of Contents

| Artikuj / Articles | Nr./No.2 | Faqe/Page |
|---|----------|-----------|
| Saktësia e analizës cefalometrike në diagnozën dhe planin e trajtimit ortodontik të pacientëve me malokluzion dento-skeletal, klasë III dhe klasë II. (Rishikim literature) | | 83 |
| Aurora Isufi, Renato Isufi, Vergjini Mulo, Lidija Kanurkova | | |
| Accuracy of cephalometric analyses in the diagnosis and orthodontic treatment planning of patients with dentoskeletal class III and class II malocclusion (Literature review) | | 94 |
| Aurora Isufi, Renato Isufi, Vergjini Mulo, Lidija Kanurkova | | |
| Defektet e lindura të buzës dhe qiellzës dhe trajtimi i tyre në shërbimin e kirurgjisë OMF, QSUT “Nënë Tereza”, Tiranë | | 107 |
| Ramazan Ifusi, Renaldo Toska, Endrit Paparisto | | |
| Cleft lip and palate and its management in OMF surgery unit, UHC “Mother Teresa”, Tirana | | 110 |
| Ramazan Ifusi, Renaldo Toska, Endrit Paparisto | | |
| Papiloma skuamoze orale e mukozës bucale: Raportim rasti | | 113 |
| Muhamet Bajrami, Irena Stojanova, Bruno Nikolovski, Mirjana Markovska Arsovska, Stavre Trajculeski, Vera Radojkova Nikolovska, Biljana Zafirova | | |
| Oral squamous papilloma of buccal mucosa: A case report | | 117 |
| Muhamet Bajrami, Irena Stojanova, Bruno Nikolovski, Mirjana Markovska Arsovska, Stavre Trajculeski, Vera Radojkova Nikolovska, Biljana Zafirova | | |
| Atricioni dentar; rehabilitimi total i një denticioni me atricion: Raportim rasti | | 120 |
| Irida Cenolli, Endrit Paparisto, Edit Xhajanka | | |
| Dental attrition; full mouth rehabilitation of dental attrition: Case report | | 125 |
| Irida Cenolli, Endrit Paparisto, Edit Xhajanka | | |
| Trajtimi endodontik i dhëmbëve të përhershëm me apeks të paformuar (Raportim rasti) | | 129 |
| Daniela Kume, Ama Mino, Adela Alushi, Klaudja Legisi | | |
| Endodontic treatment of permanent teeth with unformed apex (Case Report) | | 132 |
| Daniela Kume, Ama Mino, Adela Alushi, Klaudja Legisi | | |

ORAL SQUAMOUS PAPILLOMA OF BUCCAL MUCOSA: A CASE REPORT

Muhamet Bajrami¹, Irena Stojanova¹, Bruno Nikolovski^{1,2}, Mirjana Markovska Arsovska¹, Stavre Trajculeski¹, Vera Radojkova Nikolovska³, Biljana Zafirova⁴

¹University Dental Clinical Center "St. Pantelejmon", Clinic for Oral surgery and Implantology, Skopje, North Macedonia

²University "Goce Delcev", Faculty of Medical Sciences, Stip, North Macedonia

³University "Ss. Cyril and Methodius, Faculty of Dentistry, Department for Periodontology and Oral pathology, Skopje, North Macedonia

⁴Faculty of Medicine, Ss Cyril and Methodius University in Skopje, Macedonia

Abstract

Oral squamous papilloma is a benign proliferation of the stratified squamous epithelium, which results in a papillary or verrucous exophytic mass, that arise from the mucosal surface. The foremost common site is the palate uvula area followed by tongue, lips and buccal mucosa. The etiology remains unknown. Treatment usually requires total excision and recurrence is rare. Here, we are reporting a case of squamous papilloma in the left side of the buccal mucosa in 50 year old male patient.

Keywords: squamous papilloma, buccal mucosa, tumor, excisional biopsy.

Introduction

Oral squamous papilloma is a common lesion of the oral mucosa. It's a benign proliferation of the stratified squamous epithelium, which ends up in a well demarcated, white or normal colored lesion, that arise from the mucosal surface. The sites of predilection for localization of the lesions include the tongue, lips and soft palate (20%), but any surface of the mouth can be affected. The intraoral papilloma typically is clinically presented as slow growing, well-demarcated lesion, usually

with normal coloured mucosa and smooth surface, sessile or pedunculated base, and hard consistency, causing difficulties in mastication and speech. The dimensions of the growth is generally smaller than 1.5 cm. We present a case of 0,9 cm of squamous papilloma arising from buccal mucosa in oral cavity.

Case presentation

A 50 years old man was referred to the University Dental Clinical Center "St. Pantelejmon", Skopje, North Macedonia. Thus, the patient reported a presence of a "ball" in his mouth approximately three months. He complained on hurting the cheek while eating. This painless ball became ulcerous, started bleeding and accompanied with pain. Nevertheless, the entire medical history revealed no systemic diseases. The left submandibular lymph nodes were non palpable and non tender. The intraoral examination revealed a soft nodule with size 1,0 x 0,5cm, located on the buccal mucosa of the left side, covered by normal and healthy soft tissue. Intraorally on palpation, it was ovoid and firm in terms of consistency (Fig.1).



Figure 1. Preoperative clinical findings

There were no other lesions in the oral cavity, nor extraorally. A provisional diagnosis of Oral squamous papilloma was set.

In the pre-surgical preparation, all the normal range of preoperative routine investigations was collected. It included: complete blood count, coagulation tests and

blood pressure examination. The patient underwent surgical excision of the lesion, performed under local anesthesia. Local infiltrative anesthesia (mepivacaine (3%) with vasoconstrictor) was injected around the lesion. This was followed by surgical incision with a scalpel blade No.15 (Fig.2).



Figure 2. Instruments used for specimen removing

After the removal of the lesion, the incision was sutured with 3-0 black silk suture(Fig.3).



Figure 3. Wound suturing

Surgical excision of the lesion was performed with a 1mm depth towards the submucosa. Based on the histopathological confirmation, we set the final diagnosis of "squamous papilloma" and the patient was invited for a recall after 10 days. The histopathological report of surgical biopsy shows elliptical submucosal lesion with 0,9 x 0,6cm width and 0,1cm

thickness. Connective tissue mass lined by stratified squamous epithelium with signs of acanthosis, parakeratosis, mild papillomatosis, moderate superficial keratosis and koilocytosis. Subepithelially, dermal connective collagen tissue is seen and also scarce chronic inflammatory substrate is found. (Fig.4)




Figure 4. a) Histopathological finding

b) Microscopic view of the specimen

Discussion

Oral squamous papilloma is a generic term that is used to include papillary and verrucous growth composed of benign epithelium and connective tissue. Some are pedunculated and others are sessile. Some are single; other are multiple or diffusely involve broad areas of the oral mucosa. The squamous papilloma is the fourth most common oral mucosal mass and forms 3-4% of all biopsied oral soft tissue lesions. It was first reported as a gingival "wart" by Tomes in 1848 and is localized,

benign, HPV-induced epithelial hyperplasia. The oral squamous papilloma is usually diagnosed in people between 20-50 years. The common sites are the palate, uvula, tongue and lips. The lesions generally measure less than 1 cm in range. The lesions are generally asymptomatic as shown in our presented case above. Squamous papillomas are traditionally divided into two types: isolated-solitary and multiple-recurring. The exact etiology is unknown but the squamous papilloma is strongly associated with human papilloma virus (HPV). (Table 1)

Table1. Oral disease and associated human papilloma virus

| Oral disease | Genotypes |
|------------------------------|-----------------------------|
| | Human papilloma virus Types |
| Verruca vulgaris | 2,4 |
| Squamous papilloma | 6,11 |
| Condylomata | 6,11 |
| Focal epithelial hyperplasia | 13,32 |
| Verrucous carcinoma | 6,11,16 |

Although many oral squamous papillomas appear to be virally induced, the infectivity of the HPV must be of a very low order. The path of transmission of the virus is unknown for oral lesions. Surgical removal is the treatment of choice by either routine excision or laser ablation. Any identifiable irritant such as ill-fitting dental appliance and rough restoration should be removed. Long-term postoperative follow-up is extremely important because of the high rate of reoccurrence of incompletely removed lesion.

Conclusion

Clinical results obtained by the use of conventional surgical technique for removing squamous papillomas in 50-year-old patient was satisfactory. The technique has proven to be simple and safe. Moreover, patient discomfort is minimal and healing was rapid and effective.

References

1. Jaju PP, Suvarna PV, Desai RS. Squamous papilloma: case report and review of literature. *Int J Oral Sci.* 2010;2(4):222-5.
2. Eduardo Vidor Vieira, Angelo José Pavan, Márcio José da Silva, Squamous papilloma: treatment in dentistry; *Journal of Surgical and Clinical Dentistry* Vol.7,n.1,pp.16-19 (Oct – Dec 2015)

3. G. Sree Vijayabala, S. Mohanavalli, C. Gunasekar, K. Meenakshisundaram, U. Rajesh Kumar, VA. Janagarathinam. Squamous Papilloma of the Buccal Mucosa with Oral Submucous Fibrosis. *J Clin Med Sciences* 2(1):006-007. DOI: 10.17352/2455-8702.000018.
4. Dos Reis HL, Rabelo PC, de Santana MR, Ferreira DC, Filho AC. Oral squamous papilloma and condyloma acuminatum as manifestations of buccal-genital infection by human papillomavirus. *Indian J Sex Transm Dis AIDS.* 2009 Jan;30(1):40-2. doi: 10.4103/0253-7184.55484. PMID: 21938114; PMCID: PMC3168039.
5. J.C. Baumgartner, H. R. Stanley, and J. L. Salomone, "Zebra VI, part 2," *Journal of Endodontics*, 1991, vol. 17, no. 4, pp. 182–185.
6. Andrade SA, Pratavieira S, Paes JF, Ribeiro MM, Bagnato VS, Varotti FP. Oral squamous papilloma: a view under clinical, fluorescence and histopathological aspects. Einstein (Sao Paulo). 2019 May 9;17(2):eRC4624. English, Portuguese. doi: 10.31744/einstein_journal/2019RC4624. PMID: 31090794.
7. Rautava J, Syrjänen S. Human papillomavirus infections in the oral mucosa. *J Am Dent Assoc.* 2011 Aug;142(8):905-14. doi: 10.14219/jada.archive.2011.0297. PMID: 21804057.

ATRICIONI DENTAR; REHABILITIMI TOTAL I NJË DENTICIONI ME ATRICION : RAPORTIM RASTI

Dr. Irida Cenolli¹, Dr. Endrit Papparisto², Prof. Asoc. Edit Xhajanka³

¹*Klinika Dentare Bluetooth*

²*Klinike Dentare Private*

³*Fakulteti i Mjekësisë Dentare*

Abstrakt

Atricioni është një dukuri që e po e ndeshim gjithmonë e më shpesh në klinikat tona, madje dhe në mosha të reja. Kjo dukuri bëhet prevalente në individë në moshë të madhe, të cilët si pasojë e sëmundjeve

kronike përdorin mjekime të ndryshme që çojnë në tharje të gojës apo regurgitacion. Por vërehet dhe në mosha më të reja, të cilët kanë ndryshime në mënyrën e ushqyerjes duke ndikuar direkt në fluksin e ulët salivar,