

MINERVA

FORENSIC MEDICINE

VOLUME 143 · No. 4 · DECEMBER 2023



EDIZIONI · MINERVA · MEDICA

REVIEW

The importance of diffuse axonal injury in the daily forensic medicine practice

Natasha DAVCEVA ¹ *, Joze BALAZIC ²

¹Institute of Forensic Medicine, Faculty of Medicine, SS. Cyril and Methodius University, Skopje, Republic of North Macedonia; ²Institute of Forensic Medicine, Faculty of Medicine, University of Ljubljana, Ljubljana, Slovenia

*Corresponding author: Davceva Natasha, Institute of Forensic Medicine, Faculty of Medicine, University SS. Cyril and Methodius of Skopje, Add. Majka Theresa No 19, 1000 Skopje, Republic of North Macedonia. E-mail: natasa.davceva@ugd.edu.mk

ABSTRACT

The knowledge about the diffuse axonal injury (DAI) as a clinicopathological entity has matured in the last 30 years. It has been defined clinically (immediate and prolonged unconsciousness leading to death or severe disability) and pathologically (the triad of DAI specific changes). In terms of its biomechanics, DAI is occurring as a result of acceleration forces of longer duration and has been fully reproduced experimentally. In the process of the diagnosing of DAI, the performance of a complete forensic neuropathological examination is essential and the immunohistochemistry method using antibodies against β -amyloid precursor protein has been proved to be highly sensitive and specific, selectively targeting the damaged axons. In this review we are pointing to the significant characteristics of DAI as a distinct clinicopathological entity that can cause severe impairment of the brain function, and in the forensic medicine setting, it can be found as the concrete cause of death. We are discussing its pathological feature, its mechanism of occurrence, and the events on a cellular level, but also the dilemmas about DAI that still exist in science: 1) regarding the strict criteria for its diagnosis; 2) regarding its biomechanical significance, which can be of a big medico-legal importance.

(Cite this article as: Davceva N, Balazic J. The importance of diffuse axonal injury in the daily forensic medicine practice. *Minerva Forensic Med* 2023;143:132-40. DOI: 10.23736/S2784-8922.23.01855-1)

KEY WORDS: Diffuse axonal injury; Brain injuries; Closed head injuries.

Diffuse axonal injury (DAI) as a distinct clinicopathological entity occurring in cases with closed head injuries has been introduced in the 1980's by Adams *et al.*¹

Although, observations about “diffuse degeneration of the white brain matter” in a series of patients with severe posttraumatic dementia, have been pointed by Sabina Strich as early as some decades before.² This type of brain damage has been described under different names, such as “shearing injury”; “diffuse damage of immediate impact type” and “diffuse white matter shearing injury.”³ With all those descriptions the authors mainly referred to the basic mechanism of the occurrence of the axonal damage, that is

“the shearing and strain” of the axonal fibers following the impact.

The idea that axons in the white brain matter are susceptible to mechanical trauma developed as a result of some most meaningful experimental studies which contributed to the several major discoveries concerning the closed head injuries.

It has been clearly shown by those studies the role of acceleration as a mechanism of occurrence of the brain injury.⁴ Furthermore, it has been experimentally shown that acceleration is even sufficient mechanism to cause injury without any direct impact to the head. Two typical acceleration-deceleration injuries that occur as a result of the acceleration to the head are Acute subdural he-

matoma (ASDH), as a result of the tearing of the bridging veins, and also the Diffuse axonal injury (DAI), as a result of the shearing and strain forces to the axonal fibers. The difference between the mechanisms of those two acceleration injuries is in the duration of acceleration forces. In fact, ASDH is caused by relatively short duration (5-10 ms) angular acceleration loading at high rates of acceleration, whereas DAI occurs most readily when the head moves coronally and it occurs when the acceleration duration is longer (20-25 ms) and the rate of acceleration lower than conditions that produce ASDH.^{5, 6}

Those new discoveries lead to the introduction of the new concept of closed head injuries. With regard to the biomechanism of their occurrence they have been grouped as contact injuries which occur as a result of the direct impact to the head and inertial (acceleration-deceleration) injuries.^{4, 7}

With regard to the severity and the final outcome of one particular closed head injury, they have been classified on focal and diffuse brain damage. It has been shown that the final outcome of one particular closed head injury does not depend upon the extent of a focal brain injury as much as it depends upon the occurrence of diffuse brain injury.^{8, 9}

Revealing the existence of the diffuse brain injuries the scientists observed that in almost 50% of patients with closed head injury leading to death and with a severe impairment of the brain function, no clinical evidence of a massive intracranial lesion can be found.^{10, 11} What is more, using the routine post-mortem medical examination it can be very difficult to define the injury unless the brain is appropriately fixed before the dissection and appropriate histological examinations are undertaken.¹²⁻¹⁴

The aforementioned facts lead to the development of new methods for the purpose of post-mortal diagnosis of the diffuse brain injuries and also the implementation of a complete forensic-neuropathological examination in the daily practice.¹²⁻¹⁴ The procedure for the complete forensic-neuropathological examination has been scientifically ascertained and affirmed in the forensic medicine and already implemented in the Recommendation No. 99 of the Council of Europe about the medico-legal autopsy rules.¹²⁻¹⁵

Exactly, the development of the new methods gave us huge knowledge about the diffuse axonal injury as a clinical-pathological entity and its incidence in the overall pathology of the head trauma. In this phase, when the only methods available for the visualization of the damaged axons were the conventional staining techniques and also the methods of impregnation with silver, the axonal damage in the white brain matter was accepted as an indicator of brain trauma. With the introduction of the immunohistochemistry in the process of diagnosing of DAI, it became clear that the axonal damage does not occur only as a result of trauma, but can be also caused by other conditions, such as hypoxia and ischemia, multiple sclerosis, HIV encephalitis, infarcts, hypoglycemia, and some authors have demonstrated the existence of a high degree of axonal damage in cases of intoxication with the opiates.¹⁶⁻²² This has been a new momentum that casted a shadow on the diagnostic relevance of the axonal damage. However, soon thereafter, prominent authors reported that there is a certain difference in the pathological finding (appearance, pattern and distribution of the damaged axons) that is indicative of the origin of the axonal damage and nowadays we are pretty clear about the definition of DAI.²³⁻²⁵

What we know about the incidence of DAI in the overall pathology of the head trauma is its almost exclusive occurrence in vehicular traffic accidents and falls of considerable height *i.e.* traumatic events with a longer duration of the acceleration forces. This has been classically outlined^{1, 5} and also observed in more recent additional studies.²⁶ Some authors have pointed to the DAI as an almost universal consequence in the fatal closed head injuries.¹⁸

Given the high rate incidence of DAI in the closed head injury pathology and regarding to the complex procedure and strict criteria of its diagnosis, in this review we are going to refer to some medico-legal issues of interest associated with the DAI phenomenon.

DAI as clinicopathological entity

Considering DAI as a clinical-pathological entity that occurs in closed head injuries, there

are clinical and also pathological criteria for its diagnosis.

The most constant clinical indicator of the occurrence of DAI is the occurrence of immediate and prolonged unconsciousness, with no lucid interval and without the presence of any massive intracranial lesion.^{1, 27} This state occurs in almost half of the patients with a severe head injury and its etiology is considered to be DAI.²⁸ This has been outlined in the early papers of Adams *et al.* where has been pointed out that these patients fall into immediate and prolonged unconsciousness and stay severely disabled or in a vegetative state until death^{1, 3, 27} but also in recent studies.^{8, 21, 29} It has been emphasized that the structural base for the vegetative state after a head injury is the diffuse damage of the white brain matter.²⁹

Exploring the clinical-pathological correlation of DAI, it has been found that DAI is not a significant contributing factor to the lethal outcome from closed head injuries in the early post injury period (24 hours), but undoubtedly is a contributing factor for the severe impairment of the brain function indicated through the status of the consciousness.³⁰

Pathologically, DAI is defined by the feature of the diffuse axonal pathology widespread and throughout white brain matter of many brain regions including the white matter bundles.^{1, 3, 9, 21, 23-25, 31-33}

Given the aforementioned clinical and pathological criteria, the diagnosis of DAI is clinicopathological i.e. DAI is considered clinicopathological entity clinically characterized by the immediate and prolonged unconsciousness after the mechanical impact to the head, typically without any lucid interval, leading to severe brain failure, vegetative state and death, and pathologically defined by the feature of the widespread and disseminated damage of the axonal fibers inside the white brain matter, including the fiber tracts and the brain stem.

The mechanism of occurrence and the pathogenesis of DAI

The mechanism of the occurrence of DAI has been fully explained so far. DAI has been reproduced experimentally and this led to a pro-

found understanding of the mechanisms causing the diffuse damage of axons in the white brain matter.

Experimental studies have shown that two major mechanisms play a key role in head injuries: (1) the contact phenomenon, and (2) acceleration and deceleration.^{7, 28} Acceleration, which is a result of the sudden movement of the head, causes the pressure gradients in the intracranial cavity, thereby initiating the forces of shearing and strain. Those inertial phenomena typically produce: 1) acute subdural hematoma (ASDH) caused by the tearing of the subdural bridging blood vessels; and 2) diffuse axonal injury (DAI) in the white brain matter produced by the strain and tearing of the axonal fibres.⁴

Furthermore, it has been observed that ASDH and DAI, as two typical acceleration injuries, differ in the duration of the acceleration forces to the head. Experimentally, it has been shown that ASDH is caused by a relatively short duration (5-10 ms) of the angular acceleration loading at high rates of acceleration, whereas DAI occurs most readily where the head moves in the coronal plane and when the acceleration duration is longer (20-25 ms) and the rate of acceleration is lower than conditions that produce ASDH.^{4, 6} The role of the angular or rotational acceleration in the occurrence of DAI, especially the one in the coronal plane, has been emphasized and recently confirmed by other authors.^{4, 34, 35} Nowadays there are some novel studies which point to the combination of the translator and angular acceleration as most responsible for the occurrence of DAI.³⁶

In real life, the circumstances for the occurrence of ASDH appear in falls where the head rapidly decelerates against a firm surface, while the conditions for producing DAI appear in vehicle traffic accidents where the impact to deformable and padded surfaces lengthens the deceleration and decreases its rate. Our own results have shown that ASDH was most frequently found in cases of simple fall, assault cases, where the blow was often associated with falling on a firm surface, and in those kinds of traffic accidents where a great part of the injury mechanism has been attributed to a fall on a firm surface, represented by cyclists.³⁷ On the other hand, ASDH was least

frequently found in drivers and not found at all in passengers, which makes it an injury not so typical of vehicle traffic accidents. Analyzing the occurrence of DAI in the same study, the results have shown the overall presence of DAI in 40% of the investigated cases with a survival time of at least 2 hours, its exclusive occurrence in drivers and its high frequency in cases of a fall from a considerable height. It was not detected in the cases of a simple fall or in assault cases.³⁷

Although it has been clearly shown experimentally that the forces of acceleration and deceleration are sufficient for producing the injury, with no direct impact to the head, we are aware that it rarely happens in the real world. Almost always the direct impact to the head (either a blow or a fall), comes first. The point of great medico-legal significance is that the forces of the initial contact do not need to be strong (they do not need to be intensive enough to cause fractures of the skull), but can still produce a serious, even lethal diffuse brain injury.^{34, 38}

The events on a cellular level and the role of the β -amyloid precursor protein

The first scientific considerations about axonal injuries indicated that the underlying pathological mechanism of traumatic axonal damage, the tearing of the axons with the subsequent retraction of the torn fiber into the so called "retraction bulb," occurs at the moment of the injury, *i.e.* the primary axotomy.^{2, 3} Yet, subsequent studies showed that axonal damage is not an immediate, but rather a delayed consequence of the impairment of axoplasmic transport. (secondary axotomy).^{39, 40} According to this theory, the initial impact on the brain causes focal perturbation in the axon, resulting in focal disruption of the axoplasmic transport and subsequent accumulation of some substances that are normal contents of an axolemma, as is the β -amyloid precursor protein (β -APP). As some axons are not permanently injured in this process, but only their function is disrupted, some authors see a "potential time-window for therapeutic intervention."⁴¹

Even primary axotomy seems to be a rare phenomenon, but can still happen at the moment of the impact and prove for that are the typically

for DAI tissue-tearing in the white brain matter "gliding" hemorrhages. The rest of the events occur in the hours or days following the injury and are attributed to the progressive disorganization of the axonal cytoskeleton. Hence, the more recent studies do not reject completely the concept of primary axotomy, but consider it as a rare phenomenon probably restricted to the highest levels of injury to the axon.^{33, 42}

The β -APP is a transmembrane glycoprotein, widely represented in the central neuron system, being a constituent part of all membranous structures. It is genetically determined by the gene located in the chromosome 21.⁴³ The β -APP is part of many normal cellular functions. In the neuron, the β -APP is synthesized in the perikaryon and then, it moves through the neuron with fast anterograde transport (100-400 mm/day). Hence, in normal circumstances β -APP is not accumulated to the degree to be detected in the tissue. However, in the case of structural axonal injury, the accumulation of β -APP occurs in the proximal and the distal axonal segment to a degree where it can be detected by means of immunohistochemistry.⁴³

Thus, it has been demonstrated that all the changes that result in the degeneration of the axon and the formation of retraction bulbs occur in a time frame of about 12 to 24 hours. That is the reason why the detection of DAI in cases with a short survival time can be difficult or impossible using the conventional staining techniques. Employing the method of routine hematoxylin and eosin (H&E) staining, the damaged axons can be seen at least 24 hours after the injury, and using the methods of impregnation with silver, they can be detected 12 to 18 hours after the injury.^{33, 44}

The introduction of the immunohistochemical technique that utilizes antibodies against β -APP, represented a watershed in the elucidation of the DAI phenomenon.^{16, 17} This method enables the visualization of damaged axons as early as 2 to 3 hours post injury. This proved to be a highly specific and an extremely sensitive method targeting selectively damaged axons.^{6, 17} The application of antibodies against β -APP made the visualization of axons even in cases of short survival time (2-3 hours) possible, unlike the conven-

tional methods. Since 1994, when Sheriff introduced the method for the first time, shorter and shorter survival time has been reported where the positive β -APP immunoreactivity has been evidenced: 3 hours, 2 hours, 1.75 hours, and 35 minutes.^{6, 19, 45, 46}

Hence, the β -APP has proved to be a very useful early marker of the axonal damage and it has become one of the most useful markers in the forensic neuropathology science.

Pathology of DAI

Soon after the initial descriptions of DAI, a grading system has been conducted.⁴⁴ It has been based on a following triad of specific pathoanatomical changes:^{1, 27, 44}

- grade I has been defined as the appearance of the diffuse axonal damage in the absence of any massive macroscopic brain lesion;
- grade II has been defined with the presence of focal lesion in the corpus callosum in addition to the widespread and diffuse axonal damage (Figure 1);
- grade III of DAI is represented by a focal lesion in the dorsolateral quadrant of the rostral brain stem, mainly in the pons, in addition to the widespread and diffuse axonal damage on a histology samples (Figure 2).

Having explored the material of 60 cases with fatal closed head injuries, 30 of which were diagnosed with DAI, we have previously reported

that a focal lesion in the corpus callosum has been detected in nine cases and a focal lesion in the dorsolateral quadrant of the rostral brain stem has been detected in 10 cases.²⁶

The diagnosis of DAI must always be confirmed by the microscopic finding of the damaged axons, the swollen “varicosity-like” axons and the torn axons in the form of retraction bulbs. The criteria for the microscopic determination of the traumatic axonal damage are discussed in detail below.

Disputable questions concerning the DAI entity

Although there has been big development in the enlightening of the DAI entity so far, there are still some disputable questions in the scientific community.

The first group of questions arise from the different criteria for diagnosing DAI: Are there certain differences which can be distinguished between axonal damage caused by ischemia and axonal damage caused by trauma?, Does the secondarily occurring ischemia mask, to a greater or lesser extent, the traumatic axonal damage?, and What can be concluded with certainty from the proper interpretation of axonal damage?

At the time when DAI was originally introduced, it was considered a pure indicator of the traumatic damage of axons.^{1, 2} After the introduction of β -APP immunohistochemistry, many



Figure 1.—Focal lesion in the corpus callosum, grade II of DAI. Case with a survival time of 6 days, where a hemorrhagic focal lesion is discovered in the splenium of the corpus callosum.



Figure 2.—Focal hemorrhagic lesion in the dorsolateral segment of the rostral brainstem. Case with a several hours' survival time, where the lesion is in the transition area between the midbrain and the pons.

studies have documented β -APP immunoreactive axons in the non-traumatic cases.¹⁸⁻²² In this new era of knowledge about the axonal injuries, it has been shown that axonal damage occurs with the same frequency in cases of brain trauma and brain ischemia as well. However, it has been emphasized in these reports that axonal damage as a local isolated phenomenon should not be equalized with the entity of diffuse axonal injury and that “simultaneous demonstration of axonal injury (AI) in the corpus callosum and pons must be regarded as evidence of DAI.”⁶

A new dilemma has occurred: is the feature of the axonal damage an indicator of DAI as a traumatic entity, or are there also situations where axonal damage is not the evidence of trauma? How to make the distinction? Is DAI a distinct clinicopathological entity of traumatic origin which must be diagnosed by strict criteria?

In order to resolve this dilemma, many prominent authors have reported and explained the differences in the pathological feature of axonal damage that is indicative of its origin.^{21, 23-25} These explanations have been altogether pointing out that in the pathological determination of DAI, the diffuse axonal damage must be found in the white brain matter of many brain regions, of which at least one located above, and one below the tentorium, especially including axonal tracts, such as the corpus callosum and the internal capsule. Furthermore, the axonal damage must be with a typical traumatic appearance, pattern and distribution. These authors also emphasize that the interpretation of the pathological findings has to be in accordance with the clinical history of the head trauma. If there is a lack of appropriate clinical ante-mortem information, only the proper and correct sampling of the brain, from the exact brain regions, can help in the determination of the trauma as a cause of the axonal damage.²¹

All the current discussions are directed towards specifying the term *axonal injury* and also the diagnostic criteria where diffuse axonal injury can be confirmed. Axonal injury (AI) is a nonspecific term referring to the axonal damage of any etiology and it does not have any specific biomechanical relevance.^{6, 25} A DAI is a distinct clinicopathological entity which is clinically defined by the occurrence of immediate

and prolonged coma leading to death or severe disability, and it is pathologically defined by the widespread and disseminated axonal damage throughout the brain, including the brainstem.²⁵

Based on the above-mentioned information, there is a difference in the appearance, the distribution and the pattern of the axonal damage that is indicative of a traumatic or ischemic etiology. In the pathological determination of DAI, in addition to the macroscopic features such as focal lesions in the corpus callosum (Figure 1) and dorsolateral quadrant of the rostral pons (Figure 2), only the detection of single or small groups of scattered and diffusely arranged β -APP positive axons, seen as “sausage-like” or “varicosity-like” swollen axons or as “retraction bulbs”, can be considered as confirmation of the traumatic etiology of axonal damage (Figure 3). The detection of circumscribed foci or a linear pattern frequently described as a “zigzag” or “Z-shaped” pattern of β -APP positive axons, which in our experience are never large and are not as neatly shaped as the diffusely arranged traumatically damaged axons, was considered a predominantly hypoxic-ischemic finding and was not considered as a proof of traumatic axonal damage (Figure 4).^{23, 47}

As another difference between ischemic and traumatic β -APP expression it has been stated




Figure 3.—Typically traumatic pattern and distribution of β -APP immunoreactive damaged axons. A strong β -APP immunoreactivity is present in the corpus callosum, in a case with an 8 days' survival time. The scattered β -APP positive axons and the clearly present retraction bulbs can be noticed as well (immunohistochemical staining using the monoclonal antibody against β -amyloid precursor protein, clone 4C11, x200).




Figure 4.—A typical ischemic appearance of the β -APP expression. It can be noticed in the case with a 3 days' survival time, on a sample from the corpus callosum. There are circumscribed foci of β -APP positive axons, which are never large and are not as neatly shaped as the traumatically damaged axons. They are frequently surrounded by the dirty background. (immunohistochemical staining using the monoclonal antibody against β -amyloid precursor protein, clone 4C11, x200).

that the detection of a granulate and a dirty background is typical of cases of hypoxia and ischemia, while the detection of well-shaped damaged axons surrounded by clean background is typical of cases of trauma.^{25, 26, 31-33}

The second group of questions is especially important from the aspect of the medico-legal relevance of DAI in the process of reconstruction of the mechanism of injury. Is DAI more characteristic of certain traumatic events? Does DAI have any specific biomechanical relevance which can be of some advantage in the forensic medicine practice? Can the presence of DAI indicate the type of traumatic event that caused the head trauma?

Generally, there are two different views in the scientific community regarding this question. The first is represented by authors who stick to the classical descriptions of DAI, and according to these, a DAI is an injury mostly found in road traffic accidents (RTA) and in cases of falling from a considerable height.^{1, 3, 4, 27, 44, 48} The second view is supported by the authors whose discoveries date after the introduction of the β -APP immunohistochemistry in diagnosing DAI, and according to them DAI can be found in cases of a simple fall, and in cases of a blow, as well.⁴⁹⁻⁵³

This aspect of DAI is of huge forensic and

medicolegal importance and that is the reason why our own results have been put in the context of this discussion.^{26, 37} Analyzing the occurrence of DAI through the prism of a traumatic event (RTA, fall or blow) it was shown the highest and almost exclusive (100%) presence of DAI in drivers and also in cases of falling from a considerable height (more than 2 meters) (60%). It was not detected in the cases of a simple fall, nor in the assault cases. Those results are in accordance with the discoveries of Adams *et al.*, according to who, DAI rarely, if ever, occurs in cases of fall, unless the fall is of a considerable height.⁴⁸ Gennarelli also discussed the exclusive occurrence of DAI in RTA as early as 1983.⁵

Sporadic or isolated cases of DAI in the simple falls *i.e.* falls from somebody's own height or in cases of assault with blows to the head, have been reported occasionally.⁴⁹⁻⁵³ However, this information should be treated with caution for two reasons: first, all of these cases have been described after the introduction of the β -APP immunohistochemistry in the diagnosis of DAI, but before the recognition of the other possible reasons for axonal damage, especially ischemia; and second, the diversity of the yielded results greatly depends on the different criteria for diagnosing DAI. Diagnosing of DAI without the use of β -APP immunohistochemistry, or, diagnosing it based on the discovery of an axonal damage in one or two brain regions, is not in accordance with the novel standards for diagnosing DAI.^{21, 43} Furthermore, there are characteristics in the appearance, pattern and distribution of damaged axons that are indicative of traumatic origin.²¹⁻²⁶

Hence, we have very few cases of DAI resulting from an assault or a simple fall documented in this way, which implicates the necessity of medico-legal precautions in the process of reconstruction of the injury mechanism.

Conclusions

The information from the reviewed literature undoubtedly shows that DAI is a distinct clinicopathological entity that solely, or as part of the more complex cranial-cerebral entity, can cause serious impairment of the brain function. Speaking in the context of the forensic medicine prac-

tice, it can be found as a concrete cause of death. That is the reason why its existence should not be overlooked in the daily practice.

Today the directions about the post mortal diagnosis of DAI are specified and defined in the forensic-neuropathological science, and have been revised many times so far. According to those directions, the two main attributes in the pathological diagnosis of DAI are: 1) the diagnosed axonal damage in the white brain matter must be diffuse; and 2) it should be widespread in many brain regions, of which at least one should be located above and one below the tentorium. The pattern and the distribution of the damaged axons must suggest its traumatic origin.

The DAI entity can be detected post mortally only by performing a detailed and complete forensic-neuropathological examination of the fixed brain tissue, a fact that has been so far implemented in the international recommendations. The method of choice for the purpose of visualization of the damaged axons is the immunohistochemistry using the antibodies against β -APP.

References

- Adams JH, Graham DI, Murray LS, Scott G. Diffuse axonal injury due to nonmissile head injury in humans: an analysis of 45 cases. *Ann Neurol* 1982;12:557–63.
- Strich SJ. Diffuse degeneration of the cerebral white matter in severe dementia following head injury. *J Neurol Neurosurg Psychiatry* 1956;19:163–85.
- Adams H, Mitchell DE, Graham DI, Doyle D. Diffuse brain damage of immediate impact type. Its relationship to 'primary brain-stem damage' in head injury. *Brain* 1977;100:489–502.
- Gennarelli TA, Thibault LE, Adams JH, Graham DI, Thompson CJ, Marcincin RP. Diffuse axonal injury and traumatic coma in the primate. *Ann Neurol* 1982;12:564–74.
- Gennarelli TA. Head injury in man and experimental animals: clinical aspects. *Acta Neurochir Suppl (Wien)* 1983;32:1–13.
- Oehmichen M, Meissner C, Schmidt V, Pedal I, König HG, Saternus KS. Axonal injury—a diagnostic tool in forensic neuropathology? A review. *Forensic Sci Int* 1998;95:67–83.
- Gennarelli TA. Mechanisms of brain injury. *J Emerg Med* 1993;11(Suppl 1):5–11.
- Lobato RD, Cordobes F, Rivas JJ. Outcome from severe head injury related to the type of intracranial lesion. A CT study. *J Neurosurg* 1983;59:762–74.
- Adams JH, Jennett B, Murray LS, Teasdale GM, Gennarelli TA, Graham DI. Neuropathological findings in disabled survivors of a head injury. *J Neurotrauma* 2011;28:701–9.
- Gennarelli TA. Types and amount of axonal injury in traumatic brain injury. *Neuropathol Appl Neurobiol* 1996;22:44–52.
- Gennarelli TA, Graham DI. Neuropathology of head injuries. *Semin Clin Neuropsychiatry* 1998;3:160–75.
- Kalimo H, Saukko P, Graham D. Neuropathological examination in forensic context. *Forensic Sci Int* 2004;146:73–81.
- Kalimo H. Forensic neuropathology: an important heading in legal medicine. *Forensic Sci Int* 2004;146:71–2.
- Davceva N, Janevska V, Ilievski B, Jovanovic R. The importance of the detail forensic-neuropathological examination in the determination of the diffuse brain injuries. *Soud Lek* 2012;57:2–6.
- Brinkmann B; Committee of Ministers. Council of Europe. Harmonization of medico-legal autopsy rules. *Int J Legal Med* 1999;113:1–14.
- Sherriff FE, Bridges LR, Sivaloganathan S. Early detection of axonal injury after human head trauma using immunocytochemistry for β -amyloid precursor protein. *Acta Neuropathol* 1994;87:55–62.
- Gentleman SM, Nash MJ, Sweeting CJ, Graham DI, Roberts GW. β -amyloid precursor protein (β APP) as a marker for axonal injury after head injury. *Neurosci Lett* 1993;160:139–44.
- Gentleman SM, Roberts GW, Gennarelli TA, Maxwell WL, Adams JH, Kerr S, *et al.* Axonal injury: a universal consequence of fatal closed head injury? *Acta Neuropathol* 1995;89:537–43.
- McKenzie KJ, McLellan DR, Gentleman SM, Maxwell WL, Gennarelli TA, Graham DI. Is β -APP a marker of axonal damage in short-surviving head injury? *Acta Neuropathol* 1996;92:608–13.
- Lambri M, Djurovic V, Kibble M, Cairns N, Al-Sarraj S. Specificity and sensitivity of betaAPP in head injury. *Clin Neuropathol* 2001;20:263–71.
- Smith C, Graham DI, Geddes JF, Whitwell HL. The interpretation of beta-APP immunoreactivity: a response to C. Neiss *et al.* *Acta Neuropathol.* 2002; 104:79. *Acta Neuropathol* 2003;106:97–8.
- Niess C, Grauel U, Toennes SW, Bratzke H. Incidence of axonal injury in human brain tissue. *Acta Neuropathol* 2002;104:79–84.
- Graham DI, Smith C, Reichard R, Leclercq PD, Gentleman SM. Trials and tribulations of using β -amyloid precursor protein immunohistochemistry to evaluate traumatic brain injury in adults. *Forensic Sci Int* 2004;146:89–96.
- Reichard RR, Smith C, Graham DI. The significance of β -APP immunoreactivity in forensic practice. *Neuropathol Appl Neurobiol* 2005;31:304–13.
- Geddes JF, Whitwell HL, Graham DI. Traumatic axonal injury: practical issues for diagnosis in medicolegal cases. *Neuropathol Appl Neurobiol* 2000;26:105–16.
- Davceva N, Janevska V, Ilievski B, Spasevska L, Popeska Z. Dilemmas concerning the diffuse axonal injury as a clinicopathological entity in forensic medical practice. *J Forensic Leg Med* 2012;19:413–8.
- Adams HJ, Corsellis JA, Duchon LW. Greenfield's neuropathology. Fourth edition. New York: John Wiley & Sons; 1984.
- DiMaio VS, DiMaio D. Forensic pathology. Second edition. New York: CRC Press; 2001.
- Adams JH, Graham DI, Jennett B. The structural basis of moderate disability after traumatic brain damage. *J Neurol Neurosurg Psychiatry* 2001;71:521–4.
- Davceva N, Sivevski A, Basheska N. Traumatic axonal injury, a clinical-pathological correlation. *J Forensic Leg Med* 2017;48:35–40.

31. Graham DI, Adams JH, Murray LS, Jennett B. Neuropathology of the vegetative state after head injury. *Neuropsychol Rehabil* 2005;15:198–213.
32. Graham DI, Maxwell WL, Adams JH, Jennett B. Novel aspects of the neuropathology of the vegetative state after blunt head injury. *Prog Brain Res* 2005;150:445–55.
33. Geddes JF, Vowles GH, Beer TW, Ellison DW. The diagnosis of diffuse axonal injury: implications for forensic practice. *Neuropathol Appl Neurobiol* 1997;23:339–47.
34. Yoganandan N, Gennarelli TA, Zhang J, Pintar FA, Takhounts E, Ridella SA. Association of contact loading in diffuse axonal injuries from motor vehicle crashes. *J Trauma* 2009;66:309–15.
35. Zhang J, Yoganandan N, Pintar FA, Gennarelli TA. Role of translational and rotational accelerations on brain strain in lateral head impact. *Biomed Sci Instrum* 2006;42:501–6.
36. Wang HC, Duan ZX, Wu FF, Xie L, Zhang H, Ma YB. A new rat model for diffuse axonal injury using a combination of linear acceleration and angular acceleration. *J Neurotrauma* 2010;27:707–19.
37. Davceva N, Janevska V, Ilievski B, Petrushevska G, Popeska Z. The occurrence of acute subdural haematoma and diffuse axonal injury as two typical acceleration injuries. *J Forensic Leg Med* 2012;19:480–4.
38. Yoganandan N, Baisden JL, Maiman DJ, Gennarelli TA, Guan Y, Pintar FA, *et al.* Severe-to-fatal head injuries in motor vehicle impacts. *Accid Anal Prev* 2010;42:1370–8.
39. Povlishock JT, Christman CW. The pathobiology of traumatically induced axonal injury in animals and humans: a review of current thoughts. *J Neurotrauma* 1995;12:555–64.
40. Povlishock JT. Traumatically induced axonal injury: pathogenesis and pathobiological implications. *Brain Pathol* 1992;2:1–12.
41. Povlishock JT, Jenkins LW. Are the pathobiological changes evoked by traumatic brain injury immediate and irreversible? *Brain Pathol* 1995;5:415–26.
42. Singh A, Lu Y, Chen C, Kallakuri S, Cavanaugh JM. A new model of traumatic axonal injury to determine the effects of strain and displacement rates. *Stapp Car Crash J* 2006;50:601–23.
43. Omalu BI. Diagnosis of traumatic diffuse axonal injury. *Am J Forensic Med Pathol* 2004;25:270–3, author reply 270–1.
44. Adams JH, Doyle D, Ford I, Gennarelli TA, Graham DI, McLellan DR. Diffuse axonal injury in head injury: definition, diagnosis and grading. *Histopathology* 1989;15:49–59.
45. Blumberg PC, Scott G, Manavis J, Wainwright H, Simpson DA, McLean AJ. Staining of amyloid precursor protein to study axonal damage in mild head injury. *Lancet* 1994;344:1055–6.
46. Hortobágyi T, Wise S, Hunt N, Cary N, Djurovic V, Fegan-Earl A, *et al.* Traumatic axonal damage in the brain can be detected using β -APP immunohistochemistry within 35 min after head injury to human adults. *Neuropathol Appl Neurobiol* 2007;33:226–37.
47. Dolinak D, Matshes E. *Medicolegal neuropathology*. New York: CRC Press; 2002.
48. Adams JH, Doyle D, Graham DI, Lawrence AE, McLellan DR. Diffuse axonal injury in head injuries caused by a fall. *Lancet* 1984;2:1420–2.
49. Imajo T, Kazee AM. Diffuse axonal injury by simple fall. *Am J Forensic Med Pathol* 1992;13:169–72.
50. Raisanen J, Hougassian DF, Moskvitch M, Lawrence C. Diffuse axonal injury in a rugby player. *Am J Forensic Med Pathol* 1999;20:70–2.
51. Imajo T, Challener RC, Roessmann U. Diffuse axonal injury by assault. *Am J Forensic Med Pathol* 1987;8:217–9.
52. Imajo T, Takeshi MD. Diffuse axonal injury: its mechanism in an assault case. *Am J Forensic Med Pathol* 1996;17:324–6.
53. Graham DI, Clark JC, Adams JH, Gennarelli TA. Diffuse axonal injury caused by assault. *J Clin Pathol* 1992;45:840–1.

Conflicts of interest

Both authors certify that there is no conflict of interest with any financial organization regarding the material discussed in the manuscript.

Authors' contributions

Both authors read and approved the final version of the manuscript.

History

Manuscript accepted: September 20, 2023. - Manuscript received: September 14, 2023.