



MACEDONIAN DENTAL REVIEW

ISSN | 1 47
2545-4757 | 2024

Macedonian Dental Review is publishing by the Faculty of Dentistry, University „Ss. Cyril and Methodius“,
Skopje, Republic of North Macedonia and Macedonian Dental Society

<http://stomfak.ukim.edu.mk/msp/>

Association between anatomical and morphological features of the dental root and periodontal disease: a comprehensive review. <i>Dollaku Shkreta B., Popovska M., Georgievska Jancheska T.</i>	1
Prevalence of orthodontic anomalies in 6 years-old children with dyslalia. <i>Georgievska Jancheska T., Tosheska-Spasova N, Stavreva N.</i>	9
Extraction of maxillary impacted canine using piezosurgery - case report. <i>Gigovska Arsova A., Peeva Petreska M., Veleska Stevkovska D., Janev E., Petrova E.</i>	15
Screw Retained Vs. Cemented Retained Fixed Prosthodontic Crowns and Bridges over Dental Implants - Literature Review -. <i>Fetahu B., Elenchevski S.</i>	19
Contemporary aspects of autotransplantation of teeth (with special reference to prf-assisted autotransplantation) –review article. <i>Gjurcheski J., Veleska-Stevkovska D.</i>	27

Macedonian Dental Review is publishing by the Faculty of Dentistry, University „Ss. Cyril and Methodius“, Skopje, Republic of North Macedonia and Macedonian dental society.

Editor in chief - Prof. **Kjiro Ivanovski**, PhD, e-mail: kiroivanovski@stomfak.ukim.edu.mk

Associate editor - Prof. **Elizabeta Gjeorgievska** PhD, Department of Pediatric and Preventive Dentistry, e-mail: egjeorgievska@stomfak.ukim.edu.mk

Secretary - **Vlatko Kokalanski**, PhD, e-mail: vkokalanski@stomfak.ukim.edu.mk

Adress - Macedonian Dental Review, str. Majka Tereza br. 43 Skopje, Republic of North Macedonia, <http://stomfak.ukim.edu.mk/msp/>

Editorial Board

Mira Jankulovska, Skopje, North Macedonia
 Sonja Apostolska, Skopje, North Macedonia
 Silvana Georgieva, Skopje, North Macedonia
 Aleksandar Grchev, Skopje, North Macedonia
 Lidija Kanurkova, Skopje, North Macedonia
 Maja Pandilova, Skopje, North Macedonia
 Vesna Stevkovska, Skopje, North Macedonia
 Boris Velichkovski, Skopje, North Macedonia
 Vasilka Rendzova, Skopje, North Macedonia
 Ivan Alajbeg, Zagreb, Croatia
 Ivan Anastasov, Sophia, Bulgaria
 Nikola Angelov, Houston, USA
 Gurhan Caglayan, Hacettepe, Turkey
 Domagoj Glavina, Zagreb, Croatia
 Peter Jevnikar, Ljubljana, Slovenia
 Hristo Kisov, Plovdiv, Bulgaria
 Darko Macan, Zagreb, Croatia
 Ljubo Marion, Ljubljana, Slovenia
 John Nicholson, London, UK
 Darije Planchak, Zagreb, Croatia
 Kristina Popova, Sophia, Bulgaria
 Sasha Stankovikj, Nish, Serbia
 Dragoslav Stamenkovikj, Belgrade, Serbia
 Zrinka Tarle, Zagreb, Croatia
 Ljiljana Tihachek, Belgrade, Serbia
 Georgi Tomov, Plovdiv, Bulgaria
 Radomir Ugrinov, Sophia, Bulgaria
 Ana Angelova-Volponi, London, UK
 Vaska Vandevska, Oslo, Norway
 Radosveta Vasileva, Sophia, Bulgaria
 Miroslav Vukadinovikj, Belgrade, Serbia
 Nikola Petrichevikj, Zagreb, Croatia

Advisory board

Aleksandar Dimkov, Skopje, North Macedonia
 Ana Sotirovska Ivkovska, Skopje, North Macedonia
 Aneta Atanasovska Stojanovska, Skopje, North Macedonia
 Antonio Kirkov, Skopje, North Macedonia
 Biljana Kapusevska, Skopje, North Macedonia

Biljana Dzipunova, Skopje, North Macedonia
 Vera Radojkova Nikolovska, Skopje, North Macedonia
 Gabriela Kurchieva Cuckova, Skopje, North Macedonia
 Goran Pancevski, Skopje, North Macedonia
 Gordana Kovacevska, Skopje, North Macedonia
 Daniela Veleska Stevkoska, Skopje, North Macedonia
 Danica Popovikj Monevska, Skopje, North Macedonia
 Edvard Janev, Skopje, North Macedonia
 Emilija Bajraktarova Valjakova, Skopje, North Macedonia
 Emilija Stefanovska, Skopje, North Macedonia
 Zlatko Georgiev, Skopje, North Macedonia
 Ilijana Muratovska, Skopje, North Macedonia
 Jadranka Bundevska Josifovska, Skopje, North Macedonia
 Julijana Nikolovska, Skopje, North Macedonia
 Katarina Dirjanska, Skopje, North Macedonia
 Kristina Mitikj, Skopje, North Macedonia
 Lidija Popovska, Skopje, North Macedonia
 Marija Peeva Petreska, Skopje, North Macedonia
 Marija Stevanovikj, Skopje, North Macedonia
 Marina Eftimoska, Skopje, North Macedonia
 Marina Kacarska, Skopje, North Macedonia
 Meri Pavleska, Skopje, North Macedonia
 Mira Jankulovska, Skopje, North Macedonia
 Natasa Toseska Spasova, Skopje, North Macedonia
 Nikola Gigovski, Skopje, North Macedonia
 Oliver Dimitrovski, Skopje, North Macedonia
 Sanja Pancevska, Skopje, North Macedonia
 Saso Elencevski, Skopje, North Macedonia
 Snezana Pesevska, Skopje, North Macedonia
 Stevica Ristoska, Skopje, North Macedonia
 Suzana Dvojakovska, Skopje, North Macedonia
 Cvetanka Bajraktarova Misevska, Skopje, North Macedonia
 Gordana Apostolova, Skopje, North Macedonia
 Irena Gavrilovikj, Skopje, North Macedonia
 Sasho Jovanovski, Skopje, North Macedonia
 Aneta Mijoska, Skopje, North Macedonia
 Natasha Stavreva, Skopje, North Macedonia

Асоцијација помеѓу анатомско-морфолошките карактеристики на коренот на забот и пародонталната болест: ревијален преглед. Долаку Шкретиа Б., Појовска М., Георгиевска Јанческа Т.	1
Распространетост на ортодонтски аномалии кај деца на 6-годишна возраст со дислалија. Георгиевска Јанческа Т., Тошеска-Спасова Н, Сиварева Н.....	9
Екстракција на импактирани максиларни канини со помош на пиезохирургија - приказ на случај. Гиџовска Арсова А., Пеева Пејиреска М., Велеска Стивковска Д., Јанев Е., Пејирова Е.....	15
Зашрафени наспроти цементираны фиксны коронки и мостови над забни импланти – литературен преглед. Феџаху Б., Еленчевски С.....	19
Современи погледи на автотрансплантација на заби (со посебен осврт на прф потпомогната авторансплантација) – литературен преглед. Ѓурчески Ј., Велеска-Стивковска Д.....	27

Македонски стоматолошки преглед го издава Стоматолошкиот факултет при Универзитетот „Св. Кирил и Методиј“ Скопје, Република Северна Македонија и Македонското стоматолошко друштво.

Одговорен уредник: Проф. д-р **Ќиро Ивановски**, e-mail: kiroivanovski@stomfak.ukim.edu.mk
 Заменик одговорен уредник: Проф. д-р **Елизабета Ѓоргиевска**, e-mail: egjorgievska@stomfak.ukim.edu.mk
 Секретар на списанието: д-р **Влатко Коколански**, e-mail: vkokolanski@stomfak.ukim.edu.mk

Адреса - Македонски стоматолошки преглед, ул. Мајка Тереза бр. 43 Скопје, Република Северна Македонија
<http://stomfak.ukim.edu.mk/msp/>

Уредувачки одбор

Мира Јанкуловска, Скопје, Северна Македонија
 Соња Апостолска, Скопје, Северна Македонија
 Силвана Георгиева, Скопје, Северна Македонија
 Александар Грчев, Скопје, Северна Македонија
 Лидија Кануркова, Скопје, Северна Македонија
 Маја Пандилова, Скопје, Северна Македонија
 Весна Стевковска, Скопје, Северна Македонија
 Борис Величковски, Скопје, Северна Македонија
 Василка Ренцова, Скопје, Северна Македонија
 Иван Алајбег, Загреб, Хрватска
 Иван Анастасов, Софија, Бугарија
 Никола Ангелов, Хјустон, САД
 Гурхан Цаглајан, Хачетене, Турција
 Домагој Главина, Загреб, Хрватска
 Петер Јевникар, Љубљана, Словенија
 Христо Кисов, Пловдив, Бугарија
 Дарко Мачан, Загреб, Хрватска
 Љубо Марион, Љубљана, Словенија
 Џон Николсон, Лондон, В. Британија
 Дарије Планчак, Загреб, Хрватска
 Кристина Попова, Софија, Бугарија
 Саша Станковиќ, Ниш, Србија
 Драгослав Стаменковиќ, Белград, Србија
 Зринка Тарле, Загреб, Хрватска
 Љилана Тихачек, Белград, Србија
 Георги Томов, Пловдив, Бугарија
 Радомир Угринов, Софија, Бугарија
 Ана Ангелова Валпони, Лондон, В. Британија
 Васка Вандевска, Осло, Норвешка
 Радосвета Василева, Софија, Бугарија
 Мирослав Вукадиновиќ, Белград, Србија
 Никола Петричевиќ, Загреб, Хрватска

Советодавен одбор

Александар Димков, Скопје, Северна Македонија
 Ана Сотировска Ивковска, Скопје, Северна Македонија
 Анета Атанасовска Стојановска, Скопје, Северна Македонија

Антонио Кирков, Скопје, Северна Македонија
 Билјана Капушевска, Скопје, Северна Македонија
 Билјана Ципунова, Скопје, Северна Македонија
 Вера Радојкова Николовска, Скопје, Северна Македонија
 Габриела Курчиева Чучкова, Скопје, Северна Македонија
 Горан Панчевски, Скопје, Северна Македонија
 Гордана Ковачевска, Скопје, Северна Македонија
 Даниела Велеска Стевковска, Скопје, Северна Македонија
 Даница Поповиќ Монеvsка, Скопје, Северна Македонија
 Едвард Јанев, Скопје, Северна Македонија
 Емилија Бајрактарова Ваљакова, Скопје, Северна Македонија
 Емилија Стефановска, Скопје, Северна Македонија
 Златко Георгиев, Скопје, Северна Македонија
 Илијана Муратовска, Скопје, Северна Македонија
 Јадранка Бундевска Јосифовска, Скопје, Северна Македонија
 Јулијана Николовска, Скопје, Северна Македонија
 Катарина Дирјанска, Скопје, Северна Македонија
 Кристина Митиќ, Скопје, Северна Македонија
 Лидија Поповска, Скопје, Северна Македонија
 Марија Пеева Петреска, Скопје, Северна Македонија
 Марија Стевановиќ, Скопје, Северна Македонија
 Марина Ефтимоска, Скопје, Северна Македонија
 Марина Кацарска, Скопје, Северна Македонија
 Мери Павлеска, Скопје, Северна Македонија
 Мира Јанкуловска, Скопје, Северна Македонија
 Наташа Тошеска Спасова, Скопје, Северна Македонија
 Никола Гиговски, Скопје, Северна Македонија
 Оливер Димитровски, Скопје, Северна Македонија
 Сања Панчевска, Скопје, Северна Македонија
 Сашо Еленчевски, Скопје, Северна Македонија
 Снежана Пешевска, Скопје, Северна Македонија
 Стевица Ристоска, Скопје, Северна Македонија
 Сузана Двојаковска Божовиќ, Скопје, Северна Македонија
 Цветанка Бајрактарова Мишевска, Скопје, Северна Македонија
 Гордана Апостолова, Скопје, Северна Македонија
 Ирена Гавриловиќ, Скопје, Северна Македонија
 Сашо Јовановски, Скопје, Северна Македонија
 Анета Мијоска, Скопје, Северна Македонија
 Наташа Ставрева, Скопје, Северна Македонија

PREVALENCE OF ORTHODONTIC ANOMALIES IN 6 YEARS-OLD CHILDREN WITH DYSLALIA

РАСПРОСТРАНЕТОСТ НА ОРТОДОНТСКИ АНОМАЛИИ КАЈ ДЕЦА НА 6-ГОДИШНА ВОЗРАСТ СО ДИСЛАЛИЈА

Georgievska Jancheska T.¹, Tosheska-Spasova N.², Stavreva N.³

¹Center for Rehabilitation of Verbal Communication Pathology, Faculty of Medicine, Ss. Cyril and Methodius University in Skopje, Republic of North Macedonia, ²Department of Orthodontics, Faculty of Dentistry - Skopje, Ss. Cyril and Methodius University in Skopje, Republic of North Macedonia, ³Department of Prosthodontics, Faculty of Dentistry - Skopje, Ss. Cyril and Methodius University in Skopje, Republic of North Macedonia

Abstract

Aim of the study: The aim of our research is to determine the frequency and distribution of orthodontic anomalies in female and male children at the age of 6 who have previously been diagnosed with dyslalia. **Methods:** The frequency and distribution of orthodontic anomalies were analyzed in a selected sample (n=586) of boys and girls at the age of 6, all with dyslalia. The performed analysis is retrospective, on data for patients recorded in an outpatient clinic for carried out diagnostic orthodontic-speech examinations in the period 2015-2019 at the Public Health Institution "Center for Rehabilitation of Verbal Communication Pathology - Skopje". The data is statistically examined, represented in tables and figures and analysed descriptively. **Results:** 15% of children with dyslalia had orthodontic anomalies of teeth and jaws. Most often, in as many as 50% of cases, it is about sagittal occlusion anomalies, followed by vertical occlusion anomalies (26.1%), anomalies in the dental arches (19.3%), etc. Among children with dyslalia, girls are more likely to have orthodontic anomalies than boys. **Conclusion:** The findings obtained from the research can serve for better planning and implementation of diagnostic-rehabilitation activities in relation to orthodontic anomalies and the formation of speech sounds, with the ultimate goal of enabling, to as many children as possible, conditions for adopting clear and distinct speech sounds. **Keywords:** dyslalia, articulation disorders, orthodontics, open bite.

Апстракт

Цел на трудот: Цел на нашето истражување е да се утврди фреквенцијата и дистрибуцијата на ортодонтски неправилности кај женски и машки деца на 6-годишна возраст кај кои претходно е дијагностицирана дислалија. **Методи:** На избран примерок (n=586) од машки и женски деца на 6-годишна возраст, сите со дислалија, анализирана е фреквенцијата и дистрибуцијата на ортодонтски аномалии. Извршената анализа е ретроспективна и тоа на податоци за пациенти заведени во амбулантски дневник за спроведени дијагностички ортодотско-логопедски прегледи во периодот 2015-2019 во ЈЗУ Центар за рехабилитација на патологија на вербалната комуникација - Скопје. Податоците се статистички обработени, табеларно и графички прикажани и дескриптивно анализирани. **Резултати:** 15% од децата со дислалија имаат ортодонтска аномалија на заби и вилицы. Најчесто, во дури 50% од случаите се работи за сагитални неправилности во оклузија, потоа следуваат вертикални неправилности во оклузија (26,1%), неправилности во деналните лакови (19,3%) итн. Во рамки на половите групи, женските деца со дислалија почесто имаат ортодонтски аномалии на заби и вилицы, отколку машките деца. **Заклучок:** Добиените наоди од истражувањето може да послужат за подобро планирање и спроведување на дијагностичко-рехабилитациски активности во однос на ортодонтските аномалии и изговорот на гласови, а со крајна цел на што поголем број деца да им се овозможат услови за усвојување на чист и јасен изговор на гласови. **Клучни зборови:** дислалија, артикулациски нарушувања, ортодонција, отворен загриз.

Introduction

Orthodontics is a specific part of dentistry which deals with the study and monitoring of the growth and development of the dentition and the accompanying anatomical

structures, from birth to dental maturity, including all preventive and curative interventions in all dental irregularities that require repositioning of teeth with functional and mechanical means, in order to establish normal occlusion and adequate facial appearance¹. Orthodontics, as a science

that deals with the study of the growth and development of the orofacial system², but also the sciences that monitor the proper articulation, have a common interest in the orofacial region, a region where one of the most important human functions, speech, takes place. Any anomalies of the structures of the orofacial region can affect the correct articulation, i.e., lead to speech problems. Orthodontic anomalies are one of the causes of incorrect articulation of certain speech sounds, i.e., especially of speech sound distortion. The orthodontic anomalies that most often lead to articulation disorders are bite anomalies, which include protrusion, progonia, crossbite, open bite and cleft lip and palate³.

The incorrect articulation of one or more sounds or dyslalia is a common occurrence among school children, and if there are also orthodontic anomalies with them, it can lead to changes in the entire child's being, i.e. hinder normal psychological development, can be the cause of psychological trauma and failure at school⁴.

Different studies show that the number of articulation disorders is higher in populations with orthodontic anomalies than in eugenic patients⁵. Also, many authors agree that people with articulation disorder tend to have a proportionately high incidence of malocclusion⁶ i.e. a misalignment or improper relationship between the teeth of the upper and lower dental arches as they approach each other as the jaws close. This does not necessarily mean that the malocclusion itself is the cause of the speech disorder. However, structural malformations of the maxillary and mandibular arches can impair the production of consonants and affect speech intelligibility⁷. The effect of malocclusion on dyslalia appears to be more relevant and frequent, and increases proportionally, depending on the severity of the malocclusion⁸.

From the abovementioned, it is clear that there is a complex relationship between speech and the position of the teeth. Harvold suggested three mechanisms by which malocclusion and speech are interrelated: there may be an occlusal or skeletal problem and coincidentally an articulating problem; there may be a genetic or metabolic disorder that affects the central nervous system, leading to poor motor control and possible malformed morphogenesis; and there may be true cause-effect where occlusal or structural anomalies could affect the articulating skills⁹. Therefore, it is of particular importance to study, from as many different aspects as possible, the conditions as well as the possible relationship between orthodontic anomalies and imprecise or incorrect articulation.

In our research, the goal is to determine the frequency and distribution of orthodontic anomalies in female and male children at the age of 6 who have previously been diagnosed with dyslalia. The obtained results should be the basis for future research to determine the possible connection of orthodontic anomalies with the incorrect articulation.

Materials and methods

A retrospective analysis of data for patients recorded in the outpatient clinic for diagnostic orthodontic-speech examination in the period 2015-2019 at the Public Health Institution "Center for Rehabilitation of Verbal Communication Pathology – Skopje" (hereinafter: the Center) was carried out. Diagnostic orthodontic-speech examinations are carried out by an expert team (specialist orthodontist; specialist in ear, nose and throat; clinical speech therapist-specialist; special educator and rehabilitator; audiometrist; social worker; psychologist) according to the adopted protocol of the Center. A sample (n=586) of male (n=389) and female (n=197) 6-year-old children, all with dyslalia, was selected from the data, in which the frequency and distribution of orthodontic anomalies was analyzed. The children have normal intellectual development, without mental retardation or impaired hearing.

The data was statistically examined, represented in tables and figures and analysed descriptively. With the descriptive method, the following parameters were analyzed: gender, type of orthodontic anomalies, other anomalies of the oral structures. The statistical analysis of the data obtained from the research was done in the statistical programs Statistica for Windows 7.0 and SPSS 17.0. Pearson Chi-square test was used to compare the analyzed variables between male and female respondents. Statistical significance was defined at the $p < 0.05$.

Results

From the analyzed sample of data for children with dyslalia (n=586), children who had orthodontic anomalies of the teeth and jaws were first identified (Table 1).

It was determined that 15% of the children had orthodontic anomalies (n=88), of which 58% are male (n=51) and 42% are female (n=37).

Considered within gender groups, 13.1% of male children and 18.8% of female children had some orthodontic anomalies. The statistical analysis confirmed the difference in the distribution of children with and without orthodontic anomalies between males and females as non-significant ($p = 0.069$).

Considered by groups of orthodontic anomalies, the results show that most often it is about sagittal occlusion anomalies (50%; n=44) (protrusion was most often registered within the group, n=19), followed by vertical occlusion anomalies (26.1%; n=23) (an open bite was most often registered within the group, n=15), anomalies in the dental arches (19.3%; n=17) (crowding in the dental arches was most often registered -mandibular crowding, n=9) and anomalies in the number of teeth (2.3%;

Table 1. Frequency and distribution of orthodontic anomalies in children with dyslalia, by gender and total

Orthodontic anomalies	Male		Female		Total		p-value
	N	(%)	n	(%)	N	(%)	
With	51	(13.1)	37	(18.8)	88	(15.0)	p=0.069
Without	338	(86.9)	160	(81.2)	498	(85.0)	
Total	389	(100.0)	197	(100.0)	586	(100.0)	

p (Pearson Chi-square test)

Table 2. Distribution of orthodontic anomalies, in children with dyslalia who have only one orthodontic anomaly, by gender and total

Orthodontic anomalies	Male			Female			Total	
	(n)	[1]	[2]	(n)	[1]	[2]	(n)	[3]
	43	55.1%		35	44.9%		78	
Sagittal occlusion anomalies	23	57.5%	53.5%	17	42.5%	48.6%	40	51.3%
Angle Class II, Division 1 Malocclusion (Protrusion)	9	50.0%	20.9%	9	50.0%	25.7%	18	23.1%
Angle Class II, Division 2 Malocclusion (Deckbiss)				2	100.0%	5.7%	2	2.6%
Angle Class III Malocclusion	5	83.3%	11.6%	1	16.7%	2.9%	6	7.7%
Underbite	9	64.3%	20.9%	5	35.7%	14.3%	14	17.9%
Vertical occlusion anomalies	11	55.0%	25.6%	9	45.0%	25.7%	20	25.6%
Open bite	8	57.1%	18.6%	6	42.9%	17.1%	14	17.9%
Deep bite	3	50.0%	7.0%	3	50.0%	8.6%	6	7.7%
Anomalies of teeth number	2	100.0%	4.7%				2	2.6%
Hypodontia	1	100.0%	2.3%				1	1.3%
Hyperodontia	1	100.0%	2.3%				1	1.3%
Anomalies in the position of individual teeth	1	100.0%	2.3%				1	1.3%
Rotation	1	100.0%	2.3%				1	1.3%
Anomalies of teeth shape	1	100.0%	2.3%				1	1.3%
Fusion	1	100.0%	2.3%				1	1.3%
Anomalies in the dental arches	5	35.7%	11.6%	9	64.3%	25.7%	14	17.9%
Crowding in the dental arches	4	36.4%	9.3%	7	63.6%	20.0%	11	14.1%
Spacing in the dental arches	1	33.3%	2.3%	2	66.7%	5.7%	3	3.8%

[1] – distribution of orthodontic anomalies between male and female children

[2] – distribution of orthodontic anomalies within the gender group

[3] – distribution of orthodontic anomalies in the total sample (n=78)

n=2). One case each with anomalies in the position of individual teeth and anomalies in the shape of the tooth was registered. Transversal occlusion anomalies were not registered.

Among the children in whom an orthodontic anomalies were determined (n=88), the orthodontic anomaly was mostly isolated, independent (88.6%; n=78), then together with other anomalies of the oral structures (10.2%; n=9), and only one child was registered as having two orthodontic anomalies at the same time.

More specifically, depending on whether the children with dyslalia have only one or two orthodontic anomalies, or in addition to the orthodontic anomalies they also have other anomalies of the oral structures, and depending on gender, the results are as follows:

a) Children with dyslalia and only one orthodontic anomaly

In the cases of children with only one orthodontic anomaly (n=78), the results for orthodontic anomalies by

group (Table 2) show that most often it is sagittal occlusion anomalies (51.3%; n=40), followed by vertical occlusion anomalies (25.6%; n=20), anomalies in the dental arches (17.9%; n=14) and anomalies in the number of teeth (2.6%; n=2). One case each with anomaly in the position of the individual teeth and anomaly in the teeth shape were registered, and transversal occlusion anomalies were not registered.

Regarding the distribution of orthodontic anomalies within gender groups, more than half of males have sagittal occlusion anomalies (53.5%), followed by male children with vertical occlusion anomalies (25.6%), and anomalies in the dental arches (11.6%) (Table 2, Male - column [2]). In female children, sagittal occlusion anomalies also dominate (48.6%), followed by equally represented vertical occlusion anomalies and anomalies in the dental arches (Table 2, Female - column [2]).

Table 3. Frequency and distribution of orthodontic anomalies in children with dyslalia who, in addition to one orthodontic anomaly, also have other anomalies of the oral structures, by gender and total

Orthodontic anomalies	Other anomalies of the oral structures	Male			Female			Total	
		(n)	[1]	[2]	(n)	[1]	[2]	(n)	[3]
		7	77.8%		2	22.2%		9	
Sagittal occlusion anomalies		3	75.0%	42.9%	1	25.0%	50.0%	4	44.4%
Angle Class II, Division 1 Malocclusion (Protrusion)	'Gothic' palate (High-arched palate)	1	100.0%	14.3%				1	11.1%
Underbite	Short lingual frenulum	2	66.7%	28.6%	1	33.3%	50.0%	3	33.3%
Vertical occlusion anomalies		2	100.0%	28.6%				2	22.2%
Open bite	Short lingual frenulum	1	100.0%	14.3%				1	11.1%
Deep bite	Short lingual frenulum	1	100.0%	14.3%				1	11.1%
Anomalies in the dental arches		2	66.7%	28.6%	1	33.3%	50.0%	3	33.3%
Spacing in the dental arches	Short lingual frenulum	2	66.7%	28.6%	1	33.3%	50.0%	3	33.3%

[1] – distribution of orthodontic anomalies and anomalies of the oral structures between male and female children

[2] – distribution of orthodontic anomalies and anomalies of the oral structures within the gender group

[3] – distribution of orthodontic anomalies and anomalies of the oral structures in the total sample (n=9)

b) Children with dyslalia, orthodontic anomaly and other anomalies of the oral structures

In cases, where in addition to one orthodontic anomaly there are also other anomalies of the oral structures (n=9), and in relation to orthodontic anomalies by group (Table 3), sagittal occlusion anomalies (44.4%; n=4), anomalies in the dental arches (33.3%; n=3) and vertical occlusion anomalies (22.2%; n=2) were most often registered, while the most common anomalies of the oral structures is a short lingual frenulum (89%; n=8), and one case of 'Gothic' palate.

Regarding the distribution of orthodontic anomalies within gender groups, 42.9% of males have sagittal occlusion anomalies, followed by males with vertical occlusion anomalies and anomalies of the dental arches with equal participation (28.6%) (Table 3, Males - column [2]). In female children, sagittal occlusion anomalies and anomalies in the dental arches are equally represented (Table 3, Females - column [2]).

c) Children with dyslalia and two orthodontic anomalies

Two orthodontic anomalies, at the same time, were registered in only one case of a male child, and it is a vertical occlusion anomaly (deep bite) and a teeth shape anomaly (fusion).

Discussion

Several studies try to show the relationship between dyslalia and some changes in the stomatognathic system. There are different viewpoints and approaches to researching the relationship between dyslalia and orthodontic anomalies, all unique in their own way. In our research, the main goal was to determine the frequency and distribution of orthodontic anomalies in female and male children at the age of 6, who have already been diagnosed with dyslalia, in order to get a clearer picture of the possible influence of orthodontic anomalies on the proper articulation. The review of the literature, i.e. researches by other authors and the results obtained in our research, partially coincided in certain aspects, while in others they were different.

According to the results of our research, on the total sample of data relating to 6-year-old children with dyslalia (n=586), it was determined that 15% of the children (n=88) have orthodontic anomalies of the teeth and jaws. Karevska and Trajkovski 10, who conducted a research on the etiological factors that lead to speech disorders, with an orthodontic examination determining that in their experimental group, which consisted of subjects with speech disorders, 60.4% had a pathological orthodontic finding. 10 This result

is not in accordance with the result of our research, but it must be taken into account that Karevska and Trajkovski 10 analyzed respondents from a wide age group, between the ages of 3 and 73, while in our research, all the respondents were 6-year-olds. Farronato et al. 8 who conducted the study, which included children aged 6 to 10 years, regarding the connection between malocclusions and dyslalia, determined that only 1/6 of children with dyslalia (n=351) do not have malocclusion at the same time. These results are different from those in our research, with a note that children of a different age group were examined, and that the examinations were conducted in the department of orthodontics and the department of speech rehabilitation in the same hospital. The data from our research were obtained from diagnostic orthodontic-speech examinations carried out at the Public Health Institution "Center for Rehabilitation of Verbal Communication Pathology – Skopje", which is visited by patients with speech disorders, so perhaps this is the reason for the small percentage of established orthodontic anomalies in children diagnosed with dyslalia. The results of the study by Vranić and Hunski 11, conducted at the SUVAG Center in Zagreb (Republic of Croatia), on a sample of 51 children aged 6 to 8 years with a diagnosis of dyslalia, showed that 64.6% of them had orthodontic anomalies. Although the respondents are of a similar age group to that of our research, it is notable that there is a large difference in the frequency of orthodontic anomalies. However, it can be pointed out that our research includes ten times more data on respondents with dyslalia (n=586) than the research by Vranić and Hunski (n=51).

In their research, Vranić and Hunski 11 also covered the aspect of the gender structure of children diagnosed with dyslalia and orthodontic anomalies, and their results show that orthodontic anomalies are more common in male respondents with articulation disorders. These results are in contrast to the results obtained in our study where 13.1% of male children and 18.8% of female children diagnosed with dyslalia had some orthodontic anomaly. However, it must be pointed out that in our research, statistical analysis confirmed the difference in the distribution of children with and without orthodontic anomalies between male and female children with dyslalia as non-significant (p=0.069).

Regarding whether children with dyslalia have only one orthodontic anomaly or two, or whether they have an orthodontic anomaly together with other anomalies of the oral structures, the possibilities for making a comparison with the same or similar research by other authors were limited. The results of our research showed that in children with a diagnosis of dyslalia and an established orthodontic anomaly (n=88), the orthodontic anomaly is mostly isolated, independent (88.6%), then together with other anomalies of the oral structures (10.2%), and only

one child registered two orthodontic anomalies at the same time. In their research, Farronato et al. 8 came to the conclusion that children with dyslalia with three or more forms of malocclusion are found much more often than those with two or only one form of malocclusion.

If we look at the orthodontic anomalies in children with dyslalia (n=88), by groups of orthodontic anomalies, the results of our research show that most often it is about sagittal occlusion anomalies (50%) (protrusion was most often registered within the group), followed by vertical occlusion anomalies (26.1%) (an open bite was most often registered within the group), anomalies in the dental arches (19.3%) (crowding in the dental arches was most often registered –mandibular crowding) and anomalies in the number of teeth (2.3%). One case with anomalies in the position of individual teeth and anomalies of teeth shape was registered. Transversal occlusion anomalies were not registered. In the previous results, it was possible to make a partial comparison with the results and conclusions of other authors' research. What was notable in other authors' results is that open bite is the most frequently mentioned key orthodontic anomaly leading to speech disorders. Namely, according to Leavy, Cisneros and LeBlanc [12], who conducted a study on 150 patients, an open bite (2 mm) was the key malocclusal factor underlying speech sound errors. The authors concluded that an open bite was the occlusal trait having the most potential to negatively impact sound production. According to the research by Didem, Neşe and Tülin 6 the most consistently reported traits between the malocclusion and speech defects are anterior open bite, increased overjet and overbite, and maxillary spacing. Also Sahad et al. 13 found a significant relationship between open bite, anterior lisp, and/or tongue thrust during the articulation of the alveolars /t/, /d/, /n/, and /l/. On the other hand, according to Stojković and Anđelski 3, the orthodontic anomalies that most often lead to articulation disorders are the anomalies of the bite, which include protrusion, proclination, crossbite, open bite and cleft lip and palate. Farronato et al. 8 reported a correlation between Class III occlusion, diastema, increase in overjet, open and deep bite, and articulation disorders.

For the sake of greater relevance of the results, it is possible to include and determine the state of the dentition (primary or mixed dentition), dental anomalies (caries, tartar, etc.), other harmful habits (for example nail biting, thumb sucking, etc.), wearing an orthodontic appliance, history of orthodontic treatment in future research. In addition, it can be investigated and studied more deeply and in detail, i.e. to determine a specific connection of an articulation disorder of a certain speech sound (for example sound /s/, /r/, /l/, etc.) with orthodontic anomalies.

Conclusions

The findings obtained from the research can serve for better planning and implementation of diagnostic-rehabilitation activities in relation to orthodontic anomalies and the formation of speech sounds, with the ultimate goal of enabling, for as many children as possible, conditions for the formation of clear and distinct speech sounds. The findings show that timely and comprehensive diagnosis of children's articulation disorders and other anomalies in the orofacial region can be of crucial importance for one of the most important human functions - speech. At the same time, the importance of a multidisciplinary approach in the diagnosis and rehabilitation of speech disorders in children is evident. There is general consensus on the need for a combined approach involving different specialists, such as speech therapists and orthodontists.

Reference

1. Marković M. i saradnici. *Ortodoncija*, Beograd-Zagreb: Medicinska knjiga; 1989.
2. Zuzhelova M. *Ortodoncija 1*, Skopje: Mariv-S; 2014.
3. Stojković N, Anđelski H. Orthodontic irregularities and disorders of articulation in preschool and school age. *Zdravstvena zaštita* 2011; 40(1):67-74. DOI: 10.5937/ZZ1101067S
4. Hunski, M. Speech disturbance (dyslalia) and orthodontic anomalies. *Acta stomatologica Croatica* 1988; 22(4):251-259. Available from: <https://hrcak.srce.hr/105093>
5. Ivičević-Desnica J, Hunski M, Horga D. Izgovorni i ortodontski poremećaji u predškolskoj dobi. *Govor* 2003; 20(1-2):147-156. Available from: <https://hrcak.srce.hr/179339>
6. Didem N, Neşe G, Tülin A. The Relationship between Speech Disorders and Orthodontic Abnormalities. *Turk J Orthod* 2011; 24(1):74-80. DOI: 10.13076/1300-3550-24-1-74
7. Najjar H. E. et al. Impact of orthodontic treatment on speech and phonetics: a review. *Int J Community Med Public Health* 2023; 10(8):2995–2999. DOI: <https://doi.org/10.18203/2394-6040.ijcmph20232397>
8. Farronato G, Giannini L, Riva R, Galbiati G, Maspero C. Correlations between malocclusions and dyslalias. *Eur J Paediatr Dent* 2012; 13(1):13-18.
9. Harvold EP. Speech articulation and oral morphology. *Am Speech Hearing Assoc* 1970; 5:69-75.
10. Karevska A, Trajkovski V. Etiology of speech disorders in "Institute for rehabilitation of hearing, speech and voice" – Skopje, *J Spec Educ Rehab* 2005; 6(1-2):27-34. DOI: 10.5281/zenodo.28992
11. Vranić Đ, Hunski M. Investigation of the interconnection of the articulation disorders and orthodontic anomalies. *Govor* 1990; 7(1):87-92. Available from: <https://hrcak.srce.hr/176318>
12. Leavy KM, Cisneros GJ, LeBlanc EM. Malocclusion and its relationship to speech sound production: Redefining the effect of malocclusal traits on sound production. *Am J Orthod Dentofacial Orthop* 2016; 150(1):116-123. DOI: 10.1016/j.ajodo.2015.12.015
13. Sahad MG, Nahás ACR, Scavone-Junior H, Jabur LB, Guedes-Pinto E. Vertical interincisal trespass assessment in children with speech disorders. *Braz Oral Res* 2008; 22(3):247-51.