

DAS HISTORISCHE HAUS DER ANATOMIE IN DER KÖLLIKERSTRASSE



... UND ES BLEIBT HISTORISCH IN DER JOSEF-SCHNEIDER-STRASSE

119<sup>th</sup> Annual Meeting / 35. Arbeitstagung

der Anatomischen Gesellschaft

24. bis 26. September 2025

am Institut für Anatomie und Zellbiologie

der Universität Würzburg



Poster 56:

Title:

Dermatoglyphic patterns in population of the R. of North Macedonia

Authors:

Elizabeta Chadikovska (Skopje, North Macedonia)  
Ace Dodevski (Skopje, North Macedonia)  
Anamarija Paunkoska (Skopje, North Macedonia)  
Jovana Kordovska (Skopje, North Macedonia)  
Julija Zhivadinovik (Skopje, North Macedonia)  
Niki Matveeva (Skopje, North Macedonia)  
Biljana Zafirova (Skopje, North Macedonia)  
Biljana Bojadzieva Stojanoska (Skopje, North Macedonia)  
Biljana Trpkovska (Skopje, North Macedonia)  
enikovska@yahoo.com

Abstract:

Dermatoglyphics is the scientific study of fingerprints from the hands and fingers, soles and toes.

Dermatoglyphic study has been conducted in order to determine and analyze individual dermatoglyphic characters in healthy individuals with Macedonian and Albanian nationality.

Some dermatoglyphic parameter on the 210 palm prints were read and classified. The prints were obtained from healthy individuals of Macedonian and Albanian ethnic origin. Palm prints were taken using Cummins and Midlo's ink method .Dermatoglyphics were classified according Henry's system.

The results in our population study showed that  $UL > Wc > We > DL > A > RL$  on both hands in all participants, radial loops are present only on the second digit, index and double loops are most present as complex patterns. Formula for present triradii is  $a > t > b > d > c$ . The values for Atd angle are equal from  $35^\circ - 55^\circ$ .Most often found in hypothenar region are distal loops.

In conclusion, a dermatoglyphic patterns on the hands and fingers in population in the R. of North Macedonia have been made. Normal variability of the dermatoglyphic patterns have been presented. The results provide new possibilities for further investigation of dermatoglyphics and their biological and genetic properties. Findings from our populational study can be compared with the results of other nationalities, as well as presenting a control group for comparison with dermatoglyphic patterns in many genetical disorders and different diseases.